

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



BJ

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A01N 43/04, C12N 15/00, 11/00, 11/18, 11/02, 11/08, A61K 9/127, 38/43, 38/54, 39/395		A1	(11) International Publication Number: WO 96/32841 (43) International Publication Date: 24 October 1996 (24.10.96)
(21) International Application Number: PCT/US96/05389 (22) International Filing Date: 18 April 1996 (18.04.96) (30) Priority Data: 424,874 19 April 1995 (19.04.95) US Not furnished 29 March 1996 (29.03.96) US (71) Applicant: BURSTEIN LABORATORIES, INC. [US/US]; 33601 Avenida Calita, San Juan Capistrano, CA 92673 (US). (72) Inventors: VIRTANEN, Jorma; 2015A Los Trancos, Irvine, CA 92715 (US). VIRTANEN, Sinikka; 2015A Los Trancos, Irvine, CA 92715 (US). (74) Agents: HALLUIN, Albert, P. et al.; Pennie & Edmonds, 1155 Avenue of the Americas, New York, NY 10036 (US).		(81) Designated States: AL, AM, AU, AZ, BB, BG, BR, BY, CA, CN, CZ, EE, FI, GE, HU, IS, JP, KG, KP, KR, KZ, LK, LR, LS, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TJ, TM, TR, TT, UA, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>	
(54) Title: NOVEL THERAPEUTIC BINDING MOLECULE COMPLEXES			
(57) Abstract <p>Structures that bind to a selected target are prepared having two or more different effector molecules joined to each other by a joining component. At least one of the effector molecules such as an antibody binds to the target and at least one of the other effector molecules such as an enzyme is therapeutic. The joining component may be a liposome, protein, dendrimer polymer or complementary nucleic acid chains. The joining component may be of sufficient length and/or flexibility to permit the therapeutic molecule to physically interact with the target at the same time as the binding molecule binds with the target. The structures may be used in disease treatment, immunoassays and sensors. For in vivo genetic manipulation of cells, the liposome joining component is filled with a plurality of genetic manipulation complexes that contain a polynucleotide having a sequence for recombination and a binding site, a DNA binding protein bound to the binding site and a motor protein conjugated to the polynucleotide. Prophospholipase may be on the surface of the liposome, and when activated cleaves the liposome to release the complex.</p>			
BEST AVAILABLE COPY			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

NOVEL THERAPEUTIC BINDING MOLECULE COMPLEXES**RELATED APPLICATIONS**

This application is related to copending U.S.

5 Applications Serial No. 08/332,514 filed October 31, 1994, and 08/424,874 filed April 19, 1995, which are hereby incorporated by reference.

1. FIELD OF THE INVENTION

10 The present invention is in the field supramolecular structures that specifically bind to a selected target.

2. BACKGROUND

Organized molecular systems are well known in biology and
15 chemistry. For example, pure molecular compounds form crystals, and surface active molecular compounds form monolayers at air-water interphase and vesicles in water. Bilayers of liposomes mimic biological membranes, and biological membranes are good examples of multimolecular
20 organized systems. Viruses, in particular, are highly organized supramolecular assemblies whose complexity surpasses any man-made assembly. Another prime example is the DNA double helix, which is the result of highly selective interaction of two complementary single strand molecules. Man
25 made, or artificial examples of supramolecular systems, include cryptates, i.e., inclusion complexes of macrocyclic receptor molecules, and interrupting two dimensional hydrogen bonded network by a large capping molecule. In these state-of-the-art examples, the structure of all participating
30 molecules are highly specific.

Jean-Marie Lehn has defined supramolecular chemistry as the chemistry beyond individual molecules, i.e., the chemistry of the intermolecular bond. Early work in supramolecular chemistry involved crown ethers and cryptates, compounds based
35 on the interaction of electron pair and ion and possibly additional ion-ion interaction (J.-M. Lehn, Angew. Chem. Int. Ed. Engl. 29 (1990) 1304-1319).

Oligobipyridines form in the presence of suitable metal cations such as copper(II) double-stranded helicates. Auxiliary groups may be attached into bipyridine units. If these groups are nucleotides they may serve as recognition sites for DNA (U. Koert, M.M. Harding and J.-M. Lehn, Nature (1990) 346:339).

Most previously described hydrogen bonded supramolecules are supramolecular polymers, i.e., periodic supramolecules composed of one or two repeating units. In principle the number of repeating units of polymeric supramolecules may be larger than two but until now nobody has used more than two repeating units. Examples of this class of supramolecules includes the chain-like supramolecule formed by co-crystallization of 1:1 mixture of 2,4,6-triaminopyrimidine and a suitable barbituric acid derivative (J.-M. Lehn, M. Mascal, A. DeCian, J. Fisher, J. Chem. Soc. Chem. Commun. (1990) 479).

Polymeric supramolecules formed from a single unit may also be used. For example, a tubular supramolecule has been formed from a single cyclic peptide (M. R. Ghadiri, J. R. Granja, R. A. Milligan, D. E. McRee and N. Khazanovich, Nature (1993) 366:324-327). These polymeric supramolecules are often simply crystals or mixed crystals in which hydrogen bonding plays a predominant role in structure maintenance. Even, if these supramolecules are stable in solution, their size is variable like that of a conventional polymer.

A step towards controlling supramolecular size and shape has been the use of capping molecules to interrupt the molecular association at the desired point (J. P. Mathias, C. T. Seto, J. A. Zerkowski and G. M. Whitesides in "Molecular Recognition: Chemical and Biochemical Problems II" (Ed. S. M. Roberts) Royal Society of Chemistry). A mixture of the isocyanurate derivative (benzCA₂) and trismelamine derivative (trisM₃) gives the supramolecule (trisM₃)₂ (benzCA₂)₃. This strategy typically produces supramolecules which have 'molecular weight' of 4-10 KDa.

No process exists today for creating large molecular assemblies of deliberately chosen molecules in which the

location of the molecules in the assembly can be selected accurately with respect to each other. Nonetheless, a dire need exists for such molecular structures since they could have numerous important medical, chemical and physical applications. These applications include, but are not limited to, supramolecular drugs, drug delivery to target organs, capture of viruses and catalysts, sensors and nanotechnological components.

Polypeptides and proteins, especially enzymes, have been attached to oligonucleotides. A peptide or protein has been used as a tag for an oligonucleotide or oligonucleotide is used as a tag for a polypeptide. Techniques such as ELISA allowed to trace enzymes easier than oligonucleotides, enzymes were used as tags for oligonucleotides. PCR provides for assays of extreme sensitivity. Oligonucleotides are often used as a tag for polypeptides or peptidomimetics, so that the fate of the polypeptide can be followed *in vitro* or *in vivo*. Synthesis methods which are used to prepare these conjugates are also useful in this invention. (D. Pollard-Knight, Technique (1990) 3:113-132).

Linear single-stranded tRNA forms branched structures because there are several complementary pieces of the sequence are suitably located. Recently, several two and three dimensional structures have been formed using this principle (Y. Zhang and N. C. Seeman, J. Am. Chem. (1994) 116:1661-1669; N. C. Seeman, J. Theor. Biol. (1982) 99:237-247.). These DNA based supramolecules have been bound together to form active structures. Because several steps are typically needed to create these molecules, the overall synthesis yield can be very low (0.1-1 %) because of these steps alone.

Branched pre-mRNA is found in cells. These molecules have highly specific structures in which adenosine is always linked to guanosine. These branched RNAs have been synthesized (T. Horn and M. S. Urdea Nucleic Acid. Res. (1989) 17:6959-6967; C. Sund, A. Földesi, S.-I. Yamakage and J. Chattopahyaya, Nucleic Acid. Res. (1991) 9-12). The synthesis of branched nucleic acids has been extended to the synthesis

of nucleic acid dendrimers (R. H. E. Hudson and M. J. Damha, J. Am. Chem. Soc. (1993) 113:2119-2124).

Oligonucleotide comb and fork structures have been used for analytical purposes (M. S. Urdea, B. Warner, J. A.

5 Running, J. A. Kolberg, J. M. Clyne, R. Sanchez-Pescador and T. Horn (Chiron Corp.) PCT Int. Appl. No. WO 89/03,891 05 May 1989, U.S. Appl. No. 109,282, 15 October 1987. 112 pp).

All previously known supramolecular structures have some drawbacks. It is of interest to provide novel supramolecular
10 structures that may be adapted for a variety of uses, including disease therapy, diagnostics, assays, and electronics.

3. SUMMARY OF THE INVENTION

15 The present invention provides several different binding molecule-multienzyme complexes capable of specifically binding to a target of interest. The binding molecule-multienzyme complexes of the invention comprise two or more different effector molecules joined to each other by a joining
20 component, wherein at least one of the effector molecules has the property of binding to a molecular target, i.e. a binding effector molecule, and at least one of the other effector molecules is a therapeutic effector molecule. The joining components for use in the binding molecule-multienzyme
25 complexes of the invention may be of a variety of classes including liposomes, proteins, organic polymers (including dendrimer type polymers). Another aspect of the invention to provide binding molecule-multienzyme complexes in which the joining component is of sufficient length and/or flexibility
30 to permit the therapeutic effector molecules to physically interact with the same target as binding molecule at the same time as binding effector molecule is interacting with the target.

One aspect of the invention relates to binding molecule-
35 multienzyme complexes that are supramolecules formed by at least two supramolecular component molecules. Each supramolecular component molecule comprises at least one

effector molecule and at least one nucleic acid chain. At least one of the nucleic acid chains on at least one component molecule of the supramolecules of the invention are complementary to nucleic acid chains on at least one other component, and thus are able to bind the components of the supramolecule by the formation of double stranded nucleic acid chains between the complementary chains. The present invention also provides methods of making the supramolecules of the present invention.

10 The nucleic acid chains of the supramolecules of the invention are preferably DNA, RNA and may also contain structural analogues of DNA or RNA. Effector molecules that may be used to form the supramolecules include, but are not limited to polypeptides, proteins, lipids, sugars. These
15 effector molecules may impart chemical and physical properties to the supramolecule include, hydrophobicity, hydrophilicity, electron conductivity, fluorescence, radioactivity, biological activity, cellular toxicity, catalytic activity, molecular and cellular recognition and in vivo transport selectivity.

20 Another aspect of the invention is to provide binding molecule-multienzyme complexes of the invention that may be used to treat or prevent infectious diseases, particularly viral infectious diseases. Binding molecule-multienzyme complexes suitable for the treatment and/or prevention of
25 infectious diseases comprise effector molecules that are antibodies specific for one or more antigen on a viral particle and one or more enzyme capable of catalyzing a reaction that destroys the infectivity of the virus of interest, e.g., hydrolysis of viral coat proteins or viral
30 envelope lipids.

An effector molecule for use in the invention may also be a toxin, such as ricin, which will kill the cell, if the virus is internalized. Another aspect of the invention is to provide binding molecule-multienzyme complexes adapted for the
35 treatment of non-infectious diseases. Binding molecule-multienzyme complexes for the treatment of specific diseases may comprise binding effector molecules specific for certain

cells or tissues and effector molecules that serves to directly alleviate a given disease condition.

Another aspect of the invention is to provide binding molecule-multienzyme complexes that expedite the delivery of
5 polynucleotides and other macromolecules into the interior of cells. Such binding molecule-multienzyme complexes are supramolecular structures derived from two or more supramolecular components adapted for the internalization of macromolecules may comprise effector molecules that either
10 alone, or in combination with other effector molecules, on the same or different structure, that are capable of crosslinking receptors on the surface of a cell for transformation.

Another aspect of the invention is to provide binding molecule-multienzyme complexes useful for performing assays
15 for compounds of interest, particularly immunoassays. Supermolecular structures for use in assays typically comprise an effector molecule capable of specifically binding to a compound of interest and a second effector molecule that may be capable of producing a detectable signal, e.g., an enzyme, or
20 a second molecule capable of specifically binding to a compound of interest. Another aspect of the invention is to provide assays employing binding molecule-multienzyme complexes of the invention so as to provide for the detection and/or quantitation of compounds of interest.

25 Another aspect of the invention is to provide binding molecule-multienzyme complexes useful for the prevention and treatment of atherosclerosis and related cardiovascular disorders. Binding molecule-multienzyme complexes of the invention useful for the treatment of such diseases may
30 comprise an effector molecule that is an antibody specific for antigens in atherosclerotic plaque.

Another aspect of the invention is to provide method and compositions for the genetic manipulation of cells of interest. The methods and compositions may be used for in
35 vivo and in vivo genetic therapy. The compositions for use in genetic therapy are binding molecule-multienzyme complexes adapted for genetic manipulation. Preferably, such binding

molecule-multienzyme complexes adapted for genetic therapy comprise liposomes as joining molecules and further comprise a genetic manipulation complex. Genetic manipulation complexes comprise a polynucleotide for genetic manipulation, a motor
5 protein, and a plurality of site-specific DNA binding proteins. Preferably, the genetic manipulation complex comprises an enzyme with phospholipase activity.

4. BRIEF DESCRIPTION OF THE FIGURES

10 The invention will be better understood by reference to the appended Figures, of which:

Figure 1 is a schematic representation of the construction of a supramolecule constructed from two components.

15 Figure 2(A) is a schematic representation of the construction of a square planar supramolecule constructed from four components.

Figure 2(B) is a schematic representation of the construction of a square planar supramolecule constructed from
20 four components which is reinforced by diagonal double stranded nucleic acid chains.

Figure 2(C) is a schematic representation of the construction of a tetrahedral supramolecule constructed from four components.

25 Figure 3(A) is a schematic representation of the construction of an antibody-multienzyme supramolecule constructed from supramolecular subcomponents.

Figure 3(B) is a schematic representation of the construction of supramolecular subcomponents used in Figure
30 3(A) from molecules each containing one enzyme or antibody.

Figure 3(C) is a schematic representation of the construction of two supramolecules containing an antibody and two enzymes. The combination of these two supramolecules is able to degrade all lipid components of the virus.

35 Figure 4 is a schematic representation of a supramolecule subcomponent which is capable of forming a supramolecular cage

around a virus when it combines with a complementary supramolecule subcomponent.

Figure 5 is a schematic representation of the construction of a supramolecule for surrounding an icosahedral virus. Figure 5A is a schematic representation of a typical icosahedral virus. Figure 5B is a schematic representation of the supramolecule subcomponent of Figure 4 approaching the icosahedral virus. Figure 5C depicts a second, complementary supramolecule subcomponent approaching the icosahedral virus. Figure 5D depicts two complementary supramolecule subcomponents surrounding a icosahedral virus. Figure 5E depicts a icosahedral virus encased within a supramolecule.

Figure 6 is a schematic representation of how the analogous structure for the large molecule in Figure 4 can be prepared using smaller molecules.

Figure 7 is a schematic representation of molecules needed to construct the supramolecule of Figure 5.

Figure 8 is a schematic representation of supramolecular assemblies which are similar to the molecules shown in Figure 6.

Figure 9 is a schematic representation of the use of triple helices in supramolecular assemblies.

Figure 10 illustrates an example of a spacer molecule for connecting three nucleotides to an effector molecule.

Figure 11 illustrates an example of a second spacer molecule for connecting three nucleotides to an effector molecule.

Figure 12 illustrates an example of a method for cross-linking two complementary oligonucleotides at one end.

Figure 13 illustrates an example of the coupling of two derivatized peptide chains to form a branched peptide structure which can serve as a trivalent linker.

Figure 14 illustrates an example of a method of using the manipulation of protective groups on a trivalent spacer in order to use the trivalent spacer in oligonucleotide synthesis.

Figure 15A is a schematic representation of a supramolecule adapted for transformation of a nucleic acid of interest into a eukaryotic cell. Figure 15B is a schematic representation of a supramolecule adapted for transformation
5 into a cell. Figure 15C is a schematic representation of a supramolecule adapted for transforming a cell and the internalization, i.e., transformation process.

Figure 16 is a schematic diagram of proteinaceous effector molecule covalently coupled to a phospholipid which
10 may be incorporated into the binding molecule-multienzyme complexes of the invention.

Figure 17 is a schematic diagram of liposomal binding molecule-multienzyme complex of the invention adapted for use in treating atherosclerosis. The liposomal binding molecule-
15 multienzyme complex comprises antibodies an binding effector molecules and lipase, urokinase, phospholipase A₂, and cholesterol esterase as therapeutic effector molecules.

Figure 18 is a schematic diagram of liposomal binding molecule-multienzyme complex of the invention adapted for use
20 in treating HIV-1 infections and the interaction of the complex within an HIV-1 virion. Diffusion of the effector molecules in the liposome are demonstrated. The symbol "A" refers to phospholipase A₂ used as a therapeutic effector molecule. The symbol "C" refers to cholesterol esterase used
25 as a therapeutic effector molecule. The symbol "L" refers to lipase used as a therapeutic effector molecule.

5. DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to composition of matter
30 that have the property of specifically binding to selected molecular targets of interest, and directing a therapeutic agent to the binding site. The compounds of the invention are collectively referred to as binding molecule-multienzyme complexes of the invention comprise two or more different
35 effector molecules joined to each other by a joining component, wherein at least one of the effector molecules is a binding effector molecule and at least one of the effectors is

a therapeutic effector. The joining component may be any of a variety of forms including liposome, proteins, organic polymers, and the like.

One of the types of binding molecule-multienzyme complexes of the invention are supramolecules (also referred to herein as supramolecular assemblies and supramolecular constructions) that comprise at least two components, *i.e.*, supramolecular components. Each supramolecular component comprises an effector molecule and at least one nucleic acid chain covalently joined to the effector molecule. By placing complementary nucleic acid chains on different components, the components of the supramolecule may be bound together by the associative forces, *i.e.*, hydrogen bonding, between the complementary nucleic acid chains, thereby producing supramolecular constructions in which two or more effector molecules are joined to one another a double-stranded or partially double stranded nucleic acids.

The general concept of the supramolecular assembly embodiment of the invention may be better understood by reference to Figure 1 wherein supramolecular components A and B are joined by effector molecules M and N, respectively. Components A and B are bound to each other by the double stranded nucleic acid chain formed by complementary nucleic acid chains.

There is no theoretical limit to the number of supramolecular components that may be used to construct a particular supramolecular assembly. Rather, steric factors that could limit the number of components that can be used in a particular supramolecule may be avoided by proper design of the supramolecule using basic structural information that is well known to the person of ordinary skill in the art of biochemistry. Thus the invention provides for numerous compounds that are supramolecular assemblies, *i.e.*, supramolecules, comprising two or more supramolecular components of the invention. The supramolecular components of the invention comprise an effector molecule, *e.g.*, an antibody, covalently joined to at least one polynucleotide.

Two or more supramolecular components of the invention may be joined to one another by means of the nucleic acids moieties of the supramolecular components by employing nucleic acids that have regions of complementarily or partial

5 complementarily to one another. Thus two or more effector molecules may be joined to one another by double stranded or partially double stranded nucleic acids.

Any class of molecule can be used as effector molecule component of the subject binding molecule-multienzyme
10 complexes. Suitable molecules for use as the effector molecule moieties of the subject binding molecule-multienzyme complexes include, but are not limited to, sugars, peptides, lipids, polymers. The effector molecules of the binding molecule-multienzyme complexes may serve several different
15 functions within the binding molecule-multienzyme complexes. For example, the effector molecules may be used to provide a wide array of structural features to binding molecule-multienzyme complexes. In addition, the effector molecules can also provide certain chemical and physical properties to
20 the binding molecule-multienzyme complexes, which include, but are not limited to, hydrophobicity, hydrophilicity, electron conductivity, fluorescence, radioactivity, biological activity, cellular toxicity, catalytic activity, as well as molecular and cellular recognition and *in vivo* transport
25 selectivity. Effector molecules include a variety of protein type, including toxins, proteinases, receptors, ligands, lectins, antibodies, esterases, hormones, cell surface markers, etc.

Some effector molecules are referred to herein as
30 "binding effector" molecules. Binding effector molecules have the property of specifically binding to one or more molecules of interest, i.e., target molecules. A given binding effector molecule has the property of being a binding effector molecule with respect to target molecules. Effector binding molecules
35 may be either a member of a pair of specific binding molecules, e.g., an antibody-hapten pair or a receptor-ligand pair. Examples of binding effector molecules include, but are

not limited to, antibodies, integrins, adhesins, cell surface markers, T cell receptors, MHC proteins, and the like. By using a combination of two or more different binding effectors, each having a binding specificity for a different target, greater binding specificity of the binding molecule-multienzyme complexes of the invention may be achieved. Combinations of two or more different binding effector molecules may be used in embodiment of the invention that employ liposomes as joining components as well as other molecules as joining components.

Suitable binding effector molecules include molecules capable of specifically binding to the constant region of antibodies. These antibody-specific binding molecules may be antibodies. Such antibody specific antibodies are well known in the art and are commercially available from many sources, e.g. Sigma Biochemical (St. Louis, MO). Additionally, the antibody-specific binding molecules may be molecules other than antibodies; for example, macrophages and other leukocyte have Fc receptor proteins on their surfaces. These Fc receptors may be used as antibody-specific binding molecules so as to function as binding effector molecules. As endogenously produced antibodies may accumulate at the site or sites of infections and/or tumors, thereby directing binding molecule-multienzyme complexes.

Another class of effector molecules are referred to herein as "therapeutic effector molecules". Therapeutic effector molecules are either (1) biologically active molecules, including but not limited to, enzymes, drugs, prodrugs, enzymes, ligands specific for receptors, radionuclides, and toxins or (2) detectable labels, including, but not limited to fluorophores, enzymes, and radionuclides. A given therapeutic effector molecule may be both a biologically active molecule and detectable label. The following are some examples of therapeutic effector molecules that have been used to treat various diseases. These therapeutic effector molecules may readily be adapted for use in the claimed invention. Thrombolytic therapy is well

established in the treatment of acute myocardial infarction (Z. Bode, et. al., Z. Kardiol 83 (1994) 393). Proteinases which have been approved for this purpose include urokinase and streptokinase. Recombinant plasminogen activator, which
5 activates plasminogen occurring naturally in the blood, is also in common use. Proteolytic enzymes have been also used to treat disk herniation (chymopapain, L.G. Lenke, et. al., AORN J. 59 (1994) 1230), fallopian tube (chymotrypsin, chinese) and inflammation (carboxypeptidase N, M. Rybak et.
10 al. Pharmacology 16 (1978) 11). Other enzyme therapies studied include: Adenosine deaminase, DNase (cystic fibrosis), Glucosylceramidase (Gaucher's disease), Lipase (cystic fibrosis, G. Morrison et. al., Aliment Pharmacol. Ther. 6 (1992) 549; cancer B.A. Richards, J. R. Soc. Med. 81
15 (1988) 284), Lysozyme (immunostimulant; antibacterial agent, I.A. Vereshchagin and O.D. Zhuravleva, Vrach. Delo (1994) 103), Peptide hydrolases (antiviral agents, K.N. Veremeenko, Vrach. Delo (1994) 8), Superoxide dismutase, Terrilytin (immunostimulant), Ribonuclease (antibacterial, antiviral,
20 antitumor), Phospholipase A2 (autoimmune suppression, K. Mahlberg, et. al., Acta Ophthalmol. Suppl. 182 (1987) 166); arthritis, R.B. Zurier, et. al., Ann. Rheum. Dis. 32 (1973) 466), Phospholipase C (pulmonary microembolism, I.G. Jansson, et. al., J. Trauma 28 (1988) S 222.
25 The binding molecule multienzyme complexes of the invention comprise at least one binding effector molecule and at least one therapeutic effector molecule. The binding molecule-multienzyme complexes may comprise more than one different type of binding effector molecule and may contain
30 more than one molecule of each type of binding effector molecule. Similarly, the binding molecule-multienzyme complexes of the invention may comprise more than one different type of therapeutic effector molecule and may contain more than one molecule of each type of therapeutic
35 effector molecule present in the effector molecule. The specific effector molecules within a given binding molecule-multienzyme complexes may be selected so that the target

component molecule has therapeutic value for a given disease or diseases. The person of ordinary skill in the art, given the information provided herein, will readily be able to select appropriate effector molecules to treat a disease of
5 interest. For example, a target multi-component therapeutic molecule for the treatment of a viral disease may comprise an antibody specific for a viral coat protein or viral envelope lipid as a binding effector molecule and a proteinase specific for the same viral coat protein as a therapeutic effector
10 molecule.

The nucleic acid used to join the subject supramolecular components to each other in some embodiments of the invention, i.e., targeted multi-component supramolecules comprising two or more supramolecular components, are preferably between 5
15 and 100 bases in length, although nucleic acids may be significantly longer than 100 bases. The nucleic acid portion of the subject supramolecular components and supramolecular assemblies may be any of the wide variety nucleic acid molecules, either naturally occurring, e.g., RNA or DNA, or
20 synthetic analogs, e.g., phosphorothioates. The term "nucleic acids" as used herein, unless indicated otherwise, refers to both naturally occurring nucleic acids and synthetic analogs thereof. For many applications, it may be desirable to use synthetic analogs of natural nucleic acid rather than nucleic
25 acids because of certain properties specific to the analogs e.g., nuclease resistance and higher denaturation temperatures of double-stranded nucleic acids.

Detailed descriptions on the use and synthesis of nucleic acid analogs can be found, among other places, in U.S. Patent
30 No. 5,292,875 (phosphorothioates), U.S. Patent No. 5,218,103 (thiophosphoramidites), U.S. Patent No. 5,183,885 (phosphorothioates), U.S. Patent No. 5,151,510 (phosphorothioates), U.S. Patent No. 4,814,448 (phosphonates), U.S. Patent No. 4,096,210 (phosphorates) U.S. Patent No.
35 4,094,873 (phenylphosphorothioates), Ragle et al., Nuc. Acids. Res. 18(6):4751-4757 (1990) (phosphoramidates). Information on how to synthesize conventional nucleic acid can be found,

among other places, in Eckstein Oligonucleotide and Analogues: A Practical Approach Oxford University Press (1992). The complementary nucleic acids need not necessarily be entirely complementary with respect to one another. A nucleic acid of
5 a first supramolecular component may be complementary to only a portion of the nucleic acid moiety of a second supramolecular component or the complementarity may be over the entire length of the nucleic acid. Nucleic acid moieties of the subject supramolecular components may contain multiple
10 regions of complementarity to two or more nucleic acids moieties on additional supramolecular components thereby to be joined to permitting three or more supramolecular components to be joined to one another through hybridization. The complementarity (as measured by sequence homology) may be
15 either 100 percent or less. It will be appreciated by those of ordinary skill in the art that the strength of associating, as indicated by duplex nucleic acid melting point, may be modulated by controlling factors such as the degree of complementarity, the identity of the base pairs (e.g., GC rich
20 nucleic acids have a higher T_m than AT rich nucleic acids), the choice of a nucleic acid or nucleic acid analog, the length of the region of complementarity, and the like. The nucleic acid moieties of the subject supramolecular components may be linear or branched. Methods of producing branched
25 nucleic acids are known to the person skilled in the art, and example of how to make branched nucleic acid molecules can be found in PCT Publication No. WO 89/03891. The use of branched nucleic acids as the nucleic acids as the nucleic acid moieties of the subject supramolecular components is
30 particular interest because branched nucleic acid may be used to conveniently join three or more supramolecules components to one another through hybridization of the nucleic moieties. Triple and tetra helixes of nucleic acid chains can also be used in the supramolecules in order to provide other
35 structural characteristics, such as rigidity, to the supramolecule.

The length of the nucleic acid moieties as well as the position of the complementary base on the nucleic acids may be used to control the two and three dimensional shape of the supramolecule. For example, as depicted in Figure 2(A), a square supramolecule can be prepared by employing four components which each contain two nucleic acid chains of equal length. Similarly, as also depicted in Figure 2(C), a tetrahedral supramolecule can be formed using four components. As can be seen from Figures 2(A) and 2(C), a wide variety of two and three dimensional supramolecule structures may be formed using differing numbers of components and differing numbers of complementary nucleic acid chains. For example, supramolecules of the present invention may contain geometric configurations that generally resemble triangles, squares, pentagons, hexagons, heptagons, octagons, parallelograms, pyramids, tetrahedrons, cubes and cylinders. It should also be understood that these figures are merely schematic representations of supramolecular assemblies and that the supramolecule may not actually possess these geometric structures in solution or in crystalline form because of the due to the flexibility of double stranded nucleic acid chains as well as other solvation, electronic and steric factors that may be present in a given supramolecule.

With respect to each supramolecular component, the number of nucleic acid moieties that may be attached to a particular effector molecule may be varied greatly so as to produce supramolecular assemblies of the desired structure.

Supramolecular components of the invention may comprise one or more nucleic acid moieties. The total number of nucleic acid moieties that may be attached to an effector molecule is limited by steric hinderance and the number of potential attachment sites, problems which may be avoided by proper selection of the effector molecule and the nucleic acid moieties.

In another embodiment of the supramolecular components of the invention, more than one effector molecules may be joined to a single nucleic acid molecule. Such supramolecular

components comprising a plurality of effector molecules joined to a single nucleic acid molecule may be used to form supramolecular assemblies through a nucleic acid hybridization with the nucleic acid moieties of similar supramolecular

5 components or supramolecular components in which nucleic acid moieties are joined to only a single effector molecule.

In a preferred embodiment of the binding molecule-multienzyme complexes of the invention the individual effector molecules are joined to one another by a joining component (or
10 joining means) that gives the individual effector molecules within the binding molecule-multienzyme complexes operational freedom, i.e., freedom of movement sufficient to permit a therapeutic effector component to physically interact with the same target as a binding effector component of the same
15 binding molecule-multienzyme complexes when the binding effector component is bound to the target. Several different types of molecules and multi-molecular assemblies may serve as joining components in different embodiments of the invention. The embodiment of the invention in which the various effector
20 molecules are interconnected by joining components that permit the effector molecules to have the aforementioned degree of operational freedom are not limited by the examples provided herein. The person of ordinary skill in the art of chemistry and/or molecular biology will be able to devise numerous
25 joining components having the desired characteristics. For example conventional bifunctional linker molecules may be modified a person of ordinary skill in the art organic chemistry go as have sufficient length to function in the joining of effector molecules with operational freedom. In
30 one embodiment of the invention, the joining component is a liposome, i.e. the binding molecule-multienzyme complexes is a liposome comprising a target binding molecule and a therapeutic effector molecule. In another embodiment of the invention, the joining component is a peptide, i.e. binding
35 molecule-multienzyme complexes comprises a binding effector molecule and a therapeutic effector molecule joined to each other through a peptide. In another embodiment of the

invention, the joining component is a dendrimer, i.e., binding molecule-multienzyme complexes is a dendrimer comprising a binding effector molecule and a therapeutic effector molecule.

When the binding molecule-multienzyme complexes of the invention comprises a liposome. The effector molecules are covalently attached to one or more constituent lipids of the liposome. The effector molecule may be directly coupled to the lipid, or coupled through a linker. Linkers serving to couple an effector molecule to a lipid may be of a variety of compounds, including polynucleotides. The liposome may optionally comprise one or more therapeutic effector molecules in the internal compartment of the liposome as well as covalently attached to the surface of the liposome. Liposomes can be classified as small unilamellar vesicles, large unilamellar vesicles, cell-size unilamellar vesicles and multilamellar vesicles (J. M. Wigglesworth in Membrane Processes, Molecular Biology and Medical Applications (Eds. G. Benga, H. Baum and F. A. Kummerow), Springer-Verlag, New York, 1984). Methods of preparation of these liposomes include vortexing, sonication, detergent dialysis or dilution, infusion or reverse phase evaporation, fusion methods, addition of short-chain PC's, addition of fatty acids or detergents, rapid extrusion and transient increase in pH.

The simplest way to prepare liposomes is to add water onto the dry components, and mix. The power used in the mixing determines the size of the liposomes. Other possibility is to use small molecular weight detergents to solubilize the lipid-protein components and phospholipids. The detergent is dialyzed off the solution and liposomes form spontaneously. Also the pure lipid liposomes can be prepared in a controlled manner and lipid-protein conjugates may be added later in a detergent solution. Lipid part of the conjugate will penetrate into the bilayer and the proteins will be anchored. The detergent can be dialyzed away. This approach uses relatively small amount of detergent, which can be biocompatible, and need not be removed.

The lipid part of a liposome that is coupled to an effector molecule that is a protein preferably contains primary amino or thiol group in their polar head group. Accordingly, phosphatidylethanolamines can be directly used.

5 Thiol groups can be introduced into synthetic phospholipids, although it is not a part of natural lipids. In order to anchor the lipid-protein conjugate firmly into the liposome the lipophilicity of the lipid should be increased. This can be achieved in several ways. First, the length of the alkyl

10 chains in the lipid can be significantly increased (about C₃₀) compared what is normally used (about C₁₈) in liposomes. Secondly, the number of alkyl chains in one lipid molecule can be higher than two, e.g., five alkyl chain phospholipid can be conveniently prepared starting from mannitol. Alternatively,

15 several phospholipid molecules can be chemically coupled through their polar head groups into a controlled multimer, which is further conjugated with a protein. Thirdly, phospholipids that are thermophilic bacterial phospholipids, or an analog thereof can be used. These phospholipids are

20 formally like phospholipid dimers, in which the two phospholipids are connected via their alkyl tails. Either one or both alkyl tails can be connected. Thus, a monomolecular layer in a liposome may look like a bilayer formed by a normal phospholipid. The stability of the

25 liposomes formed by these 'dimeric' phospholipids is very high. Fourthly, polymerizable phospholipids can be used. The polymer is preferably biocompatible and biodegradable, preferably a polyester or a polyamide.

The lipid component of the embodiments of the subject

30 binding molecule-multienzyme complexes that are liposomes are preferably not substrates of lipolytic enzymes used as effector molecules. When lipase and phospholipase A₂ are used as an effector molecule, the lipids must be resistant against these enzymes. If ether bonds are used to connect alkyl

35 chains with glycerol or other polyalcohols, the resistance is automatically attained, because these two enzymes are ester hydrolases. Enzymes are also sensitive for the

stereochemistry of the lipid. Use of unnatural stereochemistry reduces or completely inhibits enzymatic activity. The liposomal lipid bilayer can be highly analogous to lipid part of cellular membranes. The liposomes of this invention may also comprise proteins in addition to the effector molecules. These proteins are not normal membrane proteins and their concentration in the blood is very low. In order to make these liposomes to resemble normal blood component or cell, additional proteins may be attached onto their surfaces. Possible proteins include albumin and glycophorin. Albumin is a molecular level scavenger, which removes fatty acids and lysolipids. These lipids are formed via the action of the lipolytic enzymes, which are a part of the liposome. Thus, albumin is an ideal additional component of the liposome, because it can perform a double function: camouflage and scavenging the hydrolytic products. Glycophorin is a membrane bound glycoprotein. It will make these liposomes to look like cells to some degree.

An advantage of the liposomes is that the number of effector molecules can be easily adjusted without affecting the size of the liposome. The various components may be supposed to be randomly distributed over the surface. Because of the rapid diffusion, the effector molecules which are needed to perform a certain function, are available when needed. Once binding effector molecules makes a contact with the target, other adhesion molecules will diffuse to the vicinity of the contact site and will bind to the target. Most targets have several binding sites near each other. The strong binding between binding effector molecules and their targets is preferred, but is not necessary for the liposomes containing digestive enzymes. For instance, by removing the lipid coat of the HIV-1, the protein core of the virus will dissolve and RNA is exposed. This happens when the virus is internalized by the cell. The lipid bilayer of the virus will fuse with the cellular membrane and the RNA is released. This is in contrast to influenza virus, which has an endocytosis pathway. The infected cell dismantles the influenza virus in

lysosomes using lipases and proteases. For HIV-1 proteases are not needed. Thus, proteases are not required needed to inactivate HIV-1 and similar viruses, while for influenza virus and other viruses with a similar life cycle, proteases
5 may be effective for inactivating the virus. The liposome internal compartment may also comprise an antisense oligonucleotide or other therapeutic compound such as a drug or enzyme, In some embodiments of the invention, the liposomes may comprise a DNase or RNase in the internal
10 compartment of the liposome or as therapeutic effector molecules on the surface of the liposome.

Many naturally occurring proteins may be used without any modification as effector molecules. Because in some cases these liposomes are used *in vivo* in humans, the human proteins
15 should be used to avoid unwanted immune responses. Human proteins can be produced in transgenic plants or animals. The amino acid sequence of proteins may also be altered through well-known genetic engineering techniques to produce mutated proteins having the desired biological functions of
20 corresponding naturally occurring protein, but adapted to coupling to lipid molecules. For example, addition of a cysteine residue, either through substitution or inserting, will add a free thiol group for coupling to a lipid molecule. Moreover the location of this cysteine can be deliberately
25 chosen. It should be located so that conjugation does not disturb the activity of the protein. Normally this means that the cysteine must be as far as possible from the active site.

In order to covalently couple a lipid with an effector molecule, the lipid must contain a functional group which has
30 high enough reactivity with heterobifunctional cross-linker. Phosphatidylethanol-amines (PE) have free aliphatic amino group and can be used directly for conjugation. The conjugation of phosphatidylethanolamines (PE) and serines (PS) with proteins is well known in the literature (Egger et. al.
35 Biochem. Biophys. Acta 1104 (1992) 45-54). Especially, if the protein contains a thiol group, a chemically well defined conjugate can be easily prepared. Several spacers are

commercially available to couple two molecules, which contain an amino and a thiol group. If a protein does not contain an a thiol group, an aliphatic amino group, may be used for conjugation. Thiol groups are preferred for conjugation sites
5 in proteins as compared with amino groups because of the lower abundance of thiol groups in most proteins.

Proteins can also be conjugated with lipids after the preparation of the liposomes so as to position the proteins on
10 the outer surface of the liposome. Phospholipids having unnatural stereochemistry may be prepared as described in the literature, for example see, (J.A. Virtanen et. al., Chem. Phys. Lipids 27 (1980) 185). D-Mannitol is tritylated to yield 1,6-ditrityl-D-mannitol, which is oxidized with lead
15 tetra-acetate and the product reduced with sodium borohydride to yield 1-trityl-sn-glycerol (1-TrG). Diacylglycerols and phospholipids can be prepared starting from 1-TrG by standard methods. For the liposomes described in this application ether bond is preferred at least in the primary hydroxyl
20 group, because it is resistant against lipase. In some embodiments an sn-hydroxyl ester bond may be useful because the unnatural stereochemistry makes this bond stable against phospholipase A₂. Other lipids for use in the subject liposomes include 2-oleoyl-3-triacontanyl-sn-glycero-1-
25 phosphatidyl ethanolamine and other 2-acyl-2-alkyl-PE's.

In another embodiment of the binding molecule-multienzyme complexes of the invention, in which the joining component is a peptide. The peptide may be either naturally occurring, e.g., spectrin or fibrin, or may be artificial. The peptide
30 joining component may take the form of a large fusion protein when the binding effector molecule and the therapeutic effector molecule are both proteins. One advantage of using binding molecule-multienzyme complexes that are fusion proteins is that the compound may be produced by standard
35 recombinant DNA production techniques such as those known to the person of ordinary skill in the art, such as described in Goedolel, Gene Expression Technology Methods in Enzymology,

Vol. 85 Academic Press, San Diego (1988). In other embodiments of the invention peptide joining components may be used in which the compound is not a fusion protein.

In other embodiments of the invention, the joining component may be a dendrimer. A dendrimer is a highly branched polymer. Dendrimers provide numerous sites of attachment for effector molecules at the termini of the dendrimer. Information on how to synthesize various dendrimers suitable for use in conjunction with this invention are well known to the person of ordinary skill in the art of organic chemistry. Enzymes may be conjugated with polymers to extend their half-life in the blood (R.F. Sherwood et. al., Biochem J. 164 (1977) 461); M.J. Knauf, et. al., J. Biol. Chem. 263 (1988) 15064; A. Abuchowski and F. Davis in "Enzymes as Drugs" (Eds. J. Holsenber and J. Roberts) John Wiley and Sons, New York, 1981, pp. 367-383). Description of how to join effector molecules to polymers can be found in R. Labeque, et. al., Proc. Natl Acad. Sci. USA 90 (1993) 3476 (Phospholipase A2 for hypercholesterolemia treatment). The structure of the polymer is not well defined and the coupling is even less defined. Only average number of the enzyme molecules per enzyme is known. The coupling of the enzyme occur typically via lysine residues and because most enzymes contain several lysines, the polymer can be coupled to any of these and possible to several lysines in the same enzyme.

The effector molecules can be chemically coupled with a polymer, which can be an artificial or a biopolymer. Craft polymers give the highest operational freedom. Dendrimers having chemically different branches can provide chemically well defined structure.

The binding molecule-multienzyme complexes of the invention may be produced in a variety of environments, either *in vitro* or *in vivo*. Binding molecule-multienzyme complexes may be constructed *in vitro* by mixing the various constituents under conditions. Conditions in the *in vitro* reaction mixture may be varied so as to influence the rate of binding molecule-

multienzyme complexes formation and the nature of the binding molecule-multienzyme complexes produced.

The binding molecule-multienzyme complexes of the present invention may be used in a very wide variety of applications which include, but are not limited to treatment of infectious disease, including HIV-1 infections, treatment of atherosclerosis, treatment of cancer, immunoassays, self-assembling resist materials, for electronic self-assembling nanocircuitry, catalytic clusters, sensors, supramolecular drugs, which are capable of encapsulating viruses and/or destroying viruses. Drug and enzyme targeting to cells and viruses may be enormously improved by using supramolecular assemblies of the invention comprising many similar or different monoclonal antibodies and several drug molecules, enzymes or other effector molecules.

It will be appreciated by the person of ordinary skill in the art that the therapeutic embodiments of the binding molecule-multienzyme complexes of the invention (e.g., compound for the treatment of cancer, viral infections, atherosclerosis) also include compounds in which effector molecules are joined to one another through conventional, i.e., non-polynucleotide, linkers. The use of non-polynucleotide linkers is well known the person of ordinary skill in the art and is described in, among other places, in several volumes of the series Methods in Enzymology, Academic Press, San Diego CA. Examples of such non-polynucleotide linkers include, 4-benzoylbenzoic Acid N-hydroxysuccinimide esters, 3-maleimidobenzoic acid N-hydroxysuccinimide esters, 1,4-phenyleneisothiocyanates, and the like. In those embodiments of the invention in which non-polynucleotide linkers are used to join effector molecules, it may be advantageous to administer a mixture of different effector molecule conjugates to a patient rather than a large supramolecule. In the treatment of HIV-1 infection for example, rather than administer a single supramolecule comprising (i) an anti-gp120 macromolecule, (ii) a phospholipase, and (iii) a proteinase, it may be desirable to

administer a formulation comprising (i) an anti-gp120-phospholipase conjugate and (ii) an anti-gp120-protease conjugate.

When the binding molecule-multienzyme complexes of the invention are used in vivo, the compounds are typically administered in a composition comprising a pharmaceutical carrier. A pharmaceutical carrier can be any compatible, non-toxic substance suitable for delivery of the therapeutic proteins and nucleic acids to the patient. Sterile water, alcohol, fats, waxes, and inert solids may be included in the carrier. Pharmaceutically accepted adjuvants (buffering agents, dispersing agent) may also be incorporated into the pharmaceutical composition.

The binding molecule-multienzyme complexes components of the invention may be administered to a patient in a variety of ways. Preferably, the pharmaceutical compositions may be administered parenterally, i.e., subcutaneously, intramuscularly or intravenously. Thus, this invention provides compositions for parenteral administration which comprise a solution of the human monoclonal antibody or a cocktail thereof dissolved in an acceptable carrier, preferably an aqueous carrier. A variety of aqueous carriers can be used, e.g., water, buffered water, 0.4% saline, 0.3% glycerine and the like. These solutions are sterile and generally free of particulate matter. These compositions may be sterilized by conventional, well known sterilization techniques. The compositions may contain pharmaceutically acceptable auxiliary substances as required to approximate physiological conditions such as pH adjusting and buffering agents, toxicity adjusting agents and the like, for example sodium acetate, sodium chloride, potassium chloride, calcium chloride, sodium lactate, etc. The concentration of antibody in these formulations can vary widely, e.g., from less than about 0.5%, usually at or at least about 1% to as much as 15 or 20% by weight and will be selected primarily based on fluid volumes, viscosities, etc., in accordance with the particular mode of administration selected.

Actual methods for preparing parenterally administrable compositions and adjustments necessary for administration to subjects will be known or apparent to those skilled in the art and are described in more detail in, for example, Remington's
5 Pharmaceutical Science, 15th Ed., Mack Publishing Company, Easton, Pa (1980), which is incorporated herein by reference.

An important use of some of the embodiments of supramolecules of the invention is a 2-dimensional supramolecular structures on semiconductor or other
10 electrically conductive surfaces so that desired patterns of self-assembling resist materials may be conveniently formed. Thus, the use of X-rays and electron beam lithography may be avoided when creating nanometerscale patterns on the semiconductor surfaces. This capability will make a completely
15 new order of nanoelectronics possible.

A second application of major importance will be self-assembling nanocircuitry using this technique. Pre-prepared diodes, transistors, capacitors, resistors, etc. and wires can be connected in highly selective ways to form two or three
20 dimensional electronic entities. Electronically conducting complementary polynucleotide chains may be used when electric contact of the nanocomponents is needed.

Binding molecule-multienzyme complexes, of the invention may also be used in catalytic and sensor
25 applications. Employing the binding molecule-multienzyme complexes methodology of the present invention, enzymes may be attached to a surface in an organized fashion in order to create desired sequential reaction. With regard to sensor applications, a sensor may be created that contains additional
30 biomolecules or organic molecules that give a photonic or electrical signal when a molecule of interest is the supermolecular assembly sensor.

Binding molecule-multienzyme complexes components of the invention may be used to provide novel immunoassays and
35 related assays for the detection of compounds of interest. Immunoassay technology is highly developed and well known to person of ordinary skill in the art, see, for example, Hudson,

Practical Immunology 3rd Ed. Oxford Publication (1989), and
Catty Antibodies: A Practical Approach Volumes 1 & 2 Oxford
University press (1989). Conventional immunoassays typically
employ antibodies conjugated to enzymes, and/or antibody-
5 antibody conjugates. It will be appreciated by one skilled in
the art that many embodiments of the binding molecule-
multienzyme complexes, preferably the supramolecular
assemblies may be substituted for the conventional antibody
conjugates used in conventional assays. Binding molecule-
10 multienzyme complexes of the invention useful or assay
comprise at least one member of a specific binding pair (e.g.,
an antibody, where the specific binding pair of molecules is
an antibody and antigen target) as an effector molecule
portion of a binding molecule-multienzyme complexes component.
15 Such binding molecule-multienzyme complexes supramolecules
may, for example comprise (1) two or more specific binding
pair members, e.g., antibodies, (2) an antibody and an enzyme
capable of generating a detectable signal, e.g., alkaline
phosphatase.
20 Numerous advantageous variants of conventional
immunoassays are enabled by employing the binding molecule
multi-enzyme complexes, preferably supramolecular assemblies
of invention instead of conventional antibody conjugates
because certain embodiments of the subject binding molecule-
25 multienzyme complexes may be assembled, disassembled, or
reassembled during an assay due to the ability of the double-
stranded nucleic acid moieties of the assembly to disassociate
or removal of the appropriate conditions. For example, (i) a
binding molecule-multienzyme complexes comprising antibody
30 joined by a double stranded nucleic acid molecule to an enzyme
producing a detectable signal may be bound to a target antigen
of interest, (ii) the binding molecule-multienzyme complexes
may then be disassociated so as to release the binding
molecule-multienzyme complexes comprising the enzyme effector
35 molecule (iii) the bound portion of the binding molecule-
multienzyme complexes may then be used to form a new binding
molecule-multienzyme complexes with a second antibody as an

effector molecule, thereby permitting the immobilization of a second molecule of interest at the same location as the bound binding molecule-multienzyme complexes portion. A person of ordinary skill in the art will appreciate that the properties
5 the subject binding molecule-multienzyme complexes structures permit many new and useful assay procedures to be performed.

The binding molecule-multienzyme complexes of the invention may be adapted so as to prevent or treat various infectious diseases, including HIV-1, the etiological agent of
10 AIDS. Specific infectious organisms may be targeted by creating, and administering in an effective amount of binding molecule-multienzyme complexes of the invention comprising as therapeutic effector molecules, (1) an antibody specific for molecule on the infections agent and (2) an enzyme capable of
15 catalyzing the modification of some integral structure of the infectious agent. For example, a binding molecule-multienzyme complexes adapted for the control of HIV-1 may comprise an antibody specific for one or more HIV-1 virus components, e.g., gp120, and one or more of the following enzymes (1) a
20 phospholipase A₂, (2) a lipase, (3) a cholesterol esterase. By including such enzymes in a binding molecule-multienzyme complexes, the lipid bilayer coat the infectious viral particle that may be destroyed. Additionally, binding molecule-multienzyme complexes of the invention adapted for
25 the treatment of the infectious disease may further comprise of protease capable of degrading a protein component of the infectious agent and/or a single stranded nucleic acid capable of hybridizing to a portion of the genome of the infectious organism of interest. In other embodiments of the binding
30 molecule-multienzyme complexes of the invention for the treatment/prevention of HIV-1 infections, soluble CD4 (e.g., TT4) may be used as a binding effector molecule to provide viral target specificity. Soluble CD4 (sCD4) fragment of CD4 binds to the HIV-1 virus in a manner similar to the binding of
35 whole CD4 (R. Fisher, et. al. Nature 331 (1988) 76-78). The production of sCD4 is described among other places in, P. Maddon, et. al., USP 5,126,433. In order to provide for the

covalent attachment of sCD4 to the remainder of the binding molecule-multienzyme complex, one or more cysteine residue may be added to the sCD4 (or CD4) by mutation. Preferably, the additional cysteine residues are located on side of the CD4 fragment that is distal to the HIV-1 binding site; preferred sites for introducing cysteine residue substitution mutations are at sCD4 amino acid residue positions 64, 364 or 374 or 377. sCD4 containing additional cysteine residues may be produced either by in vitro peptide synthesis techniques or by conventional recombinant DNA expression techniques.

Binding molecule-multienzyme complexes, particularly embodiments employing a liposome as a joining component, may be adapted for the treatment of atherosclerosis. LDL has a cholesterol and cholesterol ester core surrounded by a phospholipid layer. Into the spherical lipid droplet four apo B molecules are partially buried and are tetrahedrally located. The overall structure is well defined and proteins are oriented so that the lipophilic domain is inside the lipid and hydrophilic part is in the water. LDL is very sensitive and forms by vortexing in vitro similar deposits as observed in atherosclerotic plaque (J. C. Khoo, E. Miller, P. McLoughlin and D. Steinberg. 1988. Enhanced Macrophage Uptake of Low Density Lipoprotein after Self-Aggregation. Arteriosclerosis 8: 348-358). The lipid part and apolipoprotein B are separated and most of the protein is denatured and deposited. Lipids from several LDL particles aggregate and form droplets having diameters of 30-400 nm. An antibody which is specific for the denatured LDL can be produced by a person of ordinary skill in the art. The heterogeneity of the plaque can be largely circumvented by the use of liposomal binding molecule-multienzyme complexes. Furthermore, because liposome binding molecule-multienzyme complexes binding is mediated by multiple antibodies, it is much stronger and more selective than the binding by a single antibody. Up to 50-60 antibodies can bind the liposome to the plaque without preventing enzymatic reaction. When lipid carbon-carbon double bonds in LDL are oxidized to aldehydes

they form Schiff's bases with the lysines of apolipoprotein B (W. Palinski in "Cell Interactions in Atherosclerosis" (Eds. N.J. Severs and H. Robenek), CRC Press, Boca Raton, 1992, p. 1.). These structures in oxidized LDL are antigenic and may
5 used as targets for antibodies to be used as effector molecules, both binding effector molecules and therapeutic effector molecules. Because this reaction proceeds often many years in the plaque, the concentration of the antigens is high in contrary to circulating LDL particles, which are
10 metabolized fast. Significant part of the oxidation is performed by the metalloproteins secreted by macrophages. Unfortunately macrophages are located mainly close to the edge of the plaque and the plaque is not evenly oxidized. Accordingly antibody to oxidized LDL does not bind evenly to
15 the plaque. The enzymes coupled to the antibody would digest mainly the edges of the plaque thereby amplifying the effect of macrophages. The end result might be the rupture of the plaque.

Another possibility is to use an antibody that is even
20 moderately specific for plaque to place oxidizing enzymes near deposited LDL. Circulating LDL might also be oxidized to some degree, but it is metabolized fast and completely removed in a relatively short period of time. Antibody-multienzyme complex containing antibody for oxidized LDL and digestive enzymes is
25 then given to the patient. Oxidized LDL is now evenly distributed and the plaque is also digested evenly. Thus, one treatment cycle consists of series administration of both the oxidative and the digestive antibody-enzyme complexes. A chimeric human/murine antibody Z2D3 specific for human
30 atheroma has been prepared (J. Narula, C. Ditlow, F. Chen, H. W. Strauss and B.-A. Khaw. 1992. Non-invasive localization of experimental atherosclerotic lesions with murine/human chimeric antibody Z2D3 F(ab')₂. American Heart Association 66th Scientific Meeting, November 16-19, 1992) and may be used
35 as effector molecule in the binding molecule-multienzyme complexes of the invention. Fibrin is a major noncellular constituent of atherosclerotic plaque. Fibrin is insoluble

even, if the surrounding lipids are removed. Fibrin network will also hold the cells immobilized. Accordingly the digestion of fibrin is mandatory. Plasmin is natural fibrin hydrolase and is preferred also in this connection. The problem with several
5 active proteases is that they are inhibited in the blood and removed from the circulation. Obvious method to avoid fast removal is to use plasminogen, which can be activated after the liposome has attached onto the plaque. Activation can be performed using, for instance, urokinase or recombinant tissue
10 plasminogen activator. These two proteins are already clinically approved for the treatment of acute myocardial infarction. They can be used as such or targeted using separate liposomes. Using liposomes the enzymatic activity is released only if two different types of liposomes are simultaneously
15 present. The selectivity is increased enormously, because only a relatively large surface containing enough antigens can bind these two liposomes simultaneously. For instance, circulating lipoprotein particles will not be affected. Side effects are reduced, because freely floating liposome does not have
20 proteolytic activity. The principle of using inactive proenzyme can be extended to phospholipase subscript, which has also an inactive proenzyme. The use of proenzymes is not only advantageous for the clinical use, but also the production of proenzymes is much easier than the production of actual enzymes.
25 The presence of active phospholipase A₂ would be virtually impossible in the goat milk, while proenzyme is harmless. Cholesterol esters and triglycerides are most insoluble lipid components. By hydrolyzing them by cholesterol ester hydrolase and lipase will produce relatively easily soluble fatty acids
30 and monoglycerides. Cholesterol is the largest lipophilic component. Moreover it is the most difficult to degrade. Cholesterol oxidase will transform cholesterol to 4-cholester-3-one, which is relatively polar. Thus oxidized cholesterol should be removed from the plaque.
35 The principles presented in this application enable the purposeful construction of huge molecular assemblies having an exactly known chemical structure. For example, in Example 6, as

shown in figures 5A-D, describes the construction of a supramolecule for capturing virus particles which would have a molecular weight of about 4,000,000 Daltons.

Another aspect of the invention is to provide binding molecule-multienzyme complexes adapted so as to mediate the transfer of polynucleotides of interest into a host cell, i.e., transfection or transformation. Binding molecule-multienzyme complexes of the invention for cell transfection comprise effect of molecules capable of initiation the natural internalization machinery of a eukaryotic cell. Such effector molecules e.g., antibodies, are capable of binding to cell surface molecules, e.g., receptors, and preferably cross-linking the receptors when the effector molecules are components of a binding molecule-multienzyme complexes of the invention. A binding molecule-multi-enzyme complexes comprising multiple antibodies may increase chances of internalization by increasing the concentration of cross-linked cell-surface molecules. Additionally, sets of supramolecular components of the invention may be used to transform cells by employing the internalization machinery of the cell. For example, a first supramolecular component consisting of a cell surface receptor specific antibody joined to a nucleic acid moiety and a second supramolecular component consisting of a second cell surface receptor-specific antibody joined to a complementary nucleic acid moiety. By permitting the first and second supramolecular component nucleic acids to hybridize to one another after the antibody moieties have bound to a cell surface, receptor cross-linking, and hence internalization, may be achieved. Supramolecular assemblies of the invention may also comprise additional nucleic acids for internalization into a host cell of interest. Nucleic acid components of supramolecular assemblies for cell transformation may be detached from the supramolecular assembly in a variety of ways. As shown in Figure 15A, the nucleic acid may be detached through the use of restriction enzymes or other nucleases. Additionally, nucleic acid components may detach from binding molecule-multienzyme complexes through the process of nucleic acid denaturalization,

provided the nucleic acids are not covalently attached to the assembly. In another embodiment of the subject binding molecule-multienzyme complexes for transformation, effector molecules having phospholipase A₂ activity may be used to
5 introduce pores into a cell membrane. In other embodiments of the invention, the supramolecular assembly may comprise polyamines, e.g., spermine so as to mediate transformation.

The large scale solid phase synthesis (e.g., over 1 mmole) of oligonucleotides is difficult to achieve using previously
10 described synthesis methods. A significant problem with large scale synthesis is the efficient mixing of the heterogeneous system. Silica, polystyrene or other similar solid support particles (typically spherical) modified with polyethyleneoxide chains are commonly used as a support for oligonucleotide
15 synthesis. The growing oligonucleotide chains may form coils and stacking relationships, even between oligonucleotides on separate support particles, thereby creating a network that can prevent the efficient entry of reagents. The higher density of these spherical particles also makes efficient reaction mixing
20 even more difficult.

Large scale synthesis of oligonucleotide, e.g., 0.1-1 mole, is useful for the commercial scale production of supramolecules and supramolecular components of the invention. The following improvements of the current oligonucleotide synthesis procedure
25 solve the above-described problems surrounding large scale synthesis of oligonucleotides. First, acetonitrile is replaced with a solvent or solvent mixture that has a specific density of about one and that is also better able to solvate the heterocyclic bases of nucleotides than acetonitrile. Suitable
30 solvents having these desired properties include benzonitrile or a mixture of acetonitrile and dichlorobenzene. The density of these solvents is compatible with the use of polystyrene or comparable solid supports. Solid supports will float in these preferred solvents, thereby permitting mixing steps to be easily
35 performed. Another improvement over conventional oligonucleotide synthesis that may be used to effect large scale synthesis is the exposure of the reaction mixture to microwaves

during the coupling step. Microwaves increase molecular rotation and reduce unwanted polynucleotide network formation without subjecting the reaction mixture to excessive heat. An additional improvement over conventional oligonucleotide methods
5 synthesis is instead of monomeric amidites, dimeric or trimeric amidites may be used as building blocks. Even larger amidite multimers may be used to construct oligonucleotides; however, monomeric, dimeric and trimeric amidites and their combinations are preferred. Using dimers and trimers as building blocks
10 requires preparation of 16 dimer amidites and up to 64 trimer amidites separate. The use of multimeric amidites the number of couplings during automated synthesis is decreased significantly and accordingly the yield and purity is increased. The three above-described oligonucleotide improvements may be employed
15 separately or in combination with one another. A person of ordinary skill in the art will appreciate that an ideal combination of the above-described improvements will depend upon the length of the oligonucleotide described and the scale of the synthesis.

20 Another aspect of the invention is to provide methods and compositions for the in vivo genetic manipulation of cells in humans and other animals. Such genetic manipulation methods and compositions may be used to modify targeted cells so as to either express or not express a gene of interest. By either
25 expressing or not expressing a selected gene, desirable properties may be added to the targeted cells. The genetic manipulation methods of the invention comprise the step of administering an effective amount of a binding molecule-multienzyme complex of the invention adapted for targeted
30 genetic manipulation. The binding molecule-multienzyme complex adapted for targeted genetic manipulation may use a variety of different molecules as joining components, including liposomes, proteins, nucleic acids and the like. Preferably, the joining component is a liposome. The binding molecule-multienzyme
35 complex adapted for genetic manipulation comprises one or more effector molecules for targeting the binding molecule-multienzyme complex to specific cells of interest. Such

effector molecules for targeting include, but are not limited to, antibodies for cell surface markers specific to the cell of interest, ligands for receptors, on the cell surface, and the like. By using one or more targeting effector molecules, the binding molecule-multienzymes may be targeted to cells of interest. The binding molecule-multienzyme complexes adapted for genetic manipulation further comprise a genetic manipulation complex. In embodiments of the invention in which the binding molecule-multienzyme complex adapted for genetic manipulation employs a liposome as a joining molecule, the liposome is filled with a plurality of genetic manipulation complexes. Genetic manipulation complexes comprise one or more polynucleotide molecules of a sequence designed for homologous recombination into a specific chromosomal location or locations. The genetic manipulation complex further comprises a DNA motor protein. DNA motor proteins are described in, among other places Vallee and Sheetz, Science 271:1539-1544 (1996) and Gorlich and Mattaj, Science 1513-1518 (1996). DNA motor protein bind to intracellular scaffolding protein filaments such as actin or tubulin. DNA motor protein bind to the filaments and be transported along the filament to the nucleus. By following the intracellular filament proteins to the nucleus, the motor protein serves to transport the entire genetic manipulation complex to the nucleus. The genetic manipulation complex may further comprise a protein (either antibody or other ligand) for binding to one or more nuclear receptors on the surface of the nuclear membrane, thereby expediting the translocation of the genetic manipulative complex across the nuclear membrane. The genetic manipulation complex also comprises at least one DNA binding protein, wherein the DNA binding protein can specifically bind to a sequence of the polynucleotide of the genetic manipulation complex. The DNA binding protein may also bind directly to a site or sites on the chromosome of the targeted cells, thus providing for the specific targeting of the polynucleotide to the site of interest. The DNA binding protein serves to bring the polynucleotide of interest into close proximity with the targeted chromosomal site, thereby

facilitating homologous recombination at the site of interest. The polynucleotide of the genetic manipulation complex may include any of a variety of sequences useful for genetic manipulation. Such sequences include, but are not limited to, 5 protein encoding sequences, anti-sense encoding sequences, and disrupted protein encoding sequences. The polynucleotide may further comprise a polyadenylating tail (or signal) so as to reduce the possibility of destruction by endogenous nucleases.

Binding molecule-multienzyme complexes adapted for targeted 10 genetic manipulation that employ liposomes as joining components employ relatively small liposomes so as to promote cellular uptake. Generally, liposomes of interest are about 10-80nm in diameter. Optionally, the surface of the liposomes may comprise a trypsinogen and/or a phospholipase (phospholipase A₂ being 15 particularly preferred), or a similar proenzyme. The proenzyme may, upon activation, serve to facilitate the release of the genetic manipulation complex from the liposome.

The invention having been described above may be better 20 understood by reference to the following examples. The following examples are offered in order to illustrate the invention and should not be interpreted as limiting the invention

25 EXAMPLES

1. Illustration of Complementary Nucleic Acid Sequences

Table 1 provides examples of nucleic acid sequences and their complementary sequences that may be used in the present invention; the construction of complementary nucleic acids is 30 known to the person of ordinary skill in the art.

For the purpose of these examples, complementary chains of nucleic acids are depicted as an integer and that integer underlined. For example, $-(A_n-C_p)_i$ is identified as 1 in Table 1. Its complement, $-(T_n-G_p)_i$ is labelled 1. With regard to the 35 indices n, p, q and r used in Table 1, it should be understood that these indices are independent for each set of complementary nucleic acid chains.

TABLE 1

Chain	Unit	Complementary Unit	Complementary	Chain
5	1 - (A _n -C _p) i	(T _n -G _p) i-		<u>1</u>
	2 - (A _n -T _p) i	(T _n -A _p) i-	n≠p ^a	<u>2</u>
	3 - (C _n -G _p) i	(G _n -C _p) i-	n≠p ^a	<u>3</u>
	4 - (A _n -C _p -G _q) j	(T _n -G _p -C _q) j-		<u>4</u>
	5 - (A _n -G _p -C _q) j	(T _n -C _p -G _q) j-		<u>5</u>
10	6 - (A _n -C _p -T _q) j	(T _n -G _p -A _q) j-		<u>6</u>
	7 - (A _n -T _p -C _q) j	(T _n -A _p -G _q) j-		<u>7</u>
	8 - (A _n -C _p -A _q -G _r) j	(T _n -G _p -T _q -C _r) j-		<u>8</u>
	9 - (A _n -G _p -A _q -C _r) j	(T _n -C _p -T _q -G _r) j-		<u>9</u>

^a If n=p then the oligonucleotide is self-complementary and can be useful when similar units are coupled together.

Many of the examples given herein are provided in order to demonstrate the principles of the invention. Preferably, repeating units are avoided.

DNA and RNA triple helices are also well known and may be used to form the supramolecular assemblies of the invention. Triple helices may result from the association between T....A....T and C....G....C. As a result, nucleic acid changes, such as those listed in Table 2 can be used to form triple helices to bind different components of a supramolecule. One advantage in using triple helix structures is increased rigidity. This property can be utilized even after the supramolecule has been assembled. Triple helix forming oligonucleotides may be used as the nucleic acid moieties of the supramolecular components of the invention. Double helical structures, which are capable of binding to a third oligonucleotide, do so and give rigidity and shape to the supramolecule. The use of triple helices in supramolecular assemblies is demonstrated in Figure 8.

TABLE 2

	Center coil	Two outer coils
11	-A _n	T _n -

12	-G _n	C _h -
13	-(A _n -G _p) i	(T _n -C _p) i-
14	-(A _n -A _p -G _q) j	(T _n -T _p -C _q) j-
5 15	-(A _n -G _p -G _q) j	(T _n -C _p -C _q) j-

2. Construction Of A Square Planar Supramolecule

Figure 2(A) depicts the construction of a square planar supramolecule from four different components. Component A comprises effector molecule M to which is attached nucleic acid chains 1 and 2. Component B is formed by attaching nucleic acid chains 1 and 3 to effector molecule N. Component C is formed by attaching nucleic acid chains 2 and 3 to effector molecule P. Component D is formed by attaching two nucleic acid chains 3 to effector molecule Q. When components A, B and C are mixed, the complementary chains 2 and 2 of components A and C bind and the complementary chains 1 and 1 of components A and B bind. When component D is added, the 3 nucleic acid chains bind to the 3 chains of components A and C to form the square supramolecule depicted in Figure 2(A).

Figure 2(B) depicts how the square planar supramolecule can be stabilized by the addition of complementary nucleic acid chains that bind component A and C to component B to D. Since the distance between diagonally positioned effector molecules are 1.41 times the distance between effector molecules on the sides of the square supramolecule, the complementary nucleic acid chains used to bind the effector molecules diagonal to one another must be at least 1.41 times as long as the complementary nucleic acid chains binding the adjacent effector molecules in order to produce a supramolecular assembly with the desired shape.

3. Construct Of A Tetrahedral Supramolecule

Figure 2(C) depicts the construction of a tetrahedral supramolecule using four components. In order to form a tetrahedral supramolecule component A is attached to components 5 B, C and D by complementary nucleic acid chains. Similarly, components B, C and D are attached to the components by complementary nucleic acid chains.

4. Synthesis Of Components Of Supramolecules

10 A. Preparation of Nucleic Acid Chains

Several different high yield strategies for oligonucleotide synthesis have been developed, see, for example, M. J. Gait "Oligonucleotide Synthesis, a Practical Approach", IRL Press, Oxford, 1984; J. W. Engels and E. Uhlman, "Gene Synthesis", 15 Angew. Chem. Int. Ed. Engl. (1989) 28:716-724. These methods include the phosphate diester, phosphate triester, phosphite triester and phosphonate methods. Phosphite triester chemistry, which utilizes highly reactive phosphoramidites as starting materials is currently the most favored method of synthesis (R. 20 L. Letsinger, J. L. Finnan, G. A. Heavner and W. B. Lunsford, "Phosphite Coupling Procedure for Generating Internucleotide Links", J. Chem. Soc. (1975) 97:3278-3279; L. J. McBride and M. H. Caruthers, "An Investigation of Several Deoxynucleoside Phosphoramidites Useful for Synthesizing Deoxyoligonucleotides", 25 Tetrahedron Lett. (1983) 24:245-248) Oligonucleotides are most commonly prepared with automated synthesis (Beaucage, et al., Tetrahedron Lett. (1981) 22:1859-1862; U.S. Patent No. 4,458,066). All of the known methods are applicable and will provide molecular building blocks for the supramolecular 30 assembly principle described in this application.

Enzymatic methods for the production of oligonucleotides may also be used to synthesize the polynucleotide moieties of the supramolecular components of the invention. The polynucleotide moieties may also be produced in vivo and 35 subsequently cleaved into complementary single strands by heating, and separated by preparative electrophoresis or chromatography.

Short oligonucleotides may be coupled together chemically or enzymatically to obtain longer oligonucleotides, see, for example (S. A. Narang, et al., Meth. Enzymol. (1979) 68:90; U.S. Patent No. 4,356,270); N. G. Dolinnaya, N. I. Sokolova, D. T. Ashirbekova and Z. A. Shabarova, "The use of BrCN for assembling modified DNA duplexes and DNA-RNA hybrids; comparison with water soluble carbodiimide", Nucleic Acid Res. (1991) 19:3067-3072).

10 **B. Preparation of Effector Molecules**

Effector molecules, which contain aliphatic amino, dialkylamino, trialkylamino, thiol, formyl oxirane, α -halogenocarbonyl, isothiocyanato or hydroxysuccinimidyl ester groups of similar, may be coupled with suitably derivatized oligonucleotides using bifunctional spacers. Effector molecules which do not contain groups mentioned above may be activated so that they contain at least one of these groups for coupling. Groups that can be activated for coupling, include: carbon-carbon double and triple bonds, halogen, carbonyl, carboxyl and hydroxyl.

The amino acid residue sequence of proteins may altered through well-known genetic engineering techniques to as to produce non-naturally occurring proteins having the desired biological functions of a corresponding naturally occurring protein, but adapted for coupling to nucleic acid moieties. For example, addition of a cysteine residue, either through substitution or inserting, may add a free thiol group for coupling to a nucleic acid moiety.

30 **C. Attachment of Nucleic Acids to Effector Molecules**

Effector molecules may be attached to nucleic acids by numerous methods, including:

1. The molecular moiety is first attached to a solid support and is used as a linker for oligonucleotide synthesis. When oligonucleotide synthesis is completed the molecular moiety is detached from the solid support so that it remains covalently coupled with the oligonucleotide. An example of this procedure

is a Fmoc-protected polypeptide which is first synthesized on a solid support so that it has a terminal free serine hydroxyl group. The oligonucleotide synthesis is started from this hydroxyl group.

5 2. Molecular moieties other than nucleotides may be incorporated inside the oligonucleotide chain during the synthesis so as to provide functional groups for coupling to nucleic acids. For example, if these molecular moieties have at least two hydroxyl groups, one of which is free and another
10 which is protected by dimethoxytrityl group, then conventional oligonucleotide synthesis methods can be used to produce a nucleic acid that may readily be coupled to an effector molecule.

3. As a last step of the oligonucleotide synthesis a
15 molecular moiety having a suitable functional group for coupling may be attached at the end of the oligonucleotide chain. Again, if this molecular moiety has at least one hydroxyl group, it can be attached as nucleic acid monomer. This approach is already well known in the literature

20 4. A molecular moiety having a suitable functional group for coupling may be attached after the oligonucleotide synthesis is completed and part or all protecting groups have been removed. Especially molecular moieties attached using methods 1-3 can contain several functional groups which are protected by
25 orthogonal protecting groups. This allows stepwise removal of protective groups and allows regioselective attachment of new molecular moieties.

Methods of attaching enzymes to oligonucleotides that are known to the person of ordinary skill in the art may be used to
30 produce the supramolecular components and supramolecular structures of the invention. Descriptions of such techniques can be found in, for example, Jablonski et al. Nucl. Ac. Res. 14:6115-6128 (1986), Ruth DNA 3:123 (1984), Balaguer et al. Anal. Biochem. 180: 50-54 (1989), Balaguer et al. Anal. Biochem.
35 195: 105-110 (1991), Li et al. Nuc. Ac. Res. 15:5275-5287 (1987), Ghosh et al. Anal. Biochem. 178:43-51 (1989), Murakami

et al. Nuc. Ac. Res. 14:5587-5595 (1989), and Alves et al. Anal. Biochem. 189:40-50 (1988).

In order to covalently couple an oligonucleotide with a effector molecule, the oligonucleotide must contain a functional group which has a high enough reactivity to allow specific reaction at predetermined site. This functionality can be introduced into an oligonucleotide chain during normal automated synthesis, if suitable joint molecules are used. Possible functionalities include amino, dimethylamino, thiol, oxirane and other groups, which are more reactive than functional groups in nucleotides. A different approach is to use biotin-avidin chemistry or another high affinity specific non-covalent interaction. Several means of introducing these groups have already been published in the literature. See, for example, Leary, et al., Proc. Natl. Acad. Sci. USA (1983) 80:4045; Richardson and Gumpert, Nucl. Acid Res. (1983) 11:6167; Lenz and Kurz, Nucl. Acid Res. (1984) 12:3435; Meinkoth and Wahl, Anal. Biochem. (1984) 138:267; Smith, et al., Nucl. Acid Res. (1985) 13:2399, J. M. Coull, H. L. Weith and R. Bischoff, Tetrahedron Lett. (1987) 27:3991-3994; J. Haralambidis, M. Chai and G. W. Tregar, Nucleic Acid. Res. (1987) 15:4857-4876; B. C. F. Chu and L. E. Orgel, Nuc. Acid Res. (1988) 16:3671-3691. In addition to the added functionality of the oligonucleotide strand, a bifunctional spacer molecule is typically used to couple oligonucleotide and a effector molecule. Many of these spacers are well known in the literature and are commercially available.

1. Attachment of Nucleic Acids to Peptides

Peptides and peptide analogues are very commonly used as effector molecules. In order to attach oligonucleotides by normal nucleotide chemistry to a peptide, the peptide should have free hydroxyl groups. Primary hydroxyl groups are preferred. These can be implemented into a peptide by using protected ethanolamine on the carboxyl end and glycolic acid on the amino terminal, instead of an amino acid. As shown in

Figure 10, serine moieties can be used to give further attachment sites along the peptide backbone.

As shown in Figure 11, the peptide effector molecule can be branched and used as a multivalent effector structure. Several other multivalent effector structures are possible such as ethylene glycol dimer, trimer, etc. Ethylene glycol derivatives can be connected to polyalcohol to get multivalent effector structures. In order to fully exploit the present invention, conjugation of several nucleic acid chains to a single effector molecule must be possible. The process of combining nucleic acids with polymeric support and with the use of spacer molecules is well known. Similar chemistry can be used in connection with this invention to combine nucleic acid chains with effector molecules such as proteins or polypeptides.

One method for conjugating several nucleic acid chains to a single effector molecule is described below. The hydroxyl group of 2-(2'-aminoethoxy)ethanol (AAE) is first protected by t-butyldimethylsilylchloride (TBS). The product is coupled with FMOC-t-BOC-L-lysine. FMOC-group is removed and two FMOC-glycines are attached similarly. FMOC-L-glutaminic acid -t-butyl ester is the next component and will later serve as a branching point (see Figure 13). Peptide chain is extended with two glycines and one lysine. The amino group of the last lysine is reacted with propylene oxide whereby a secondary hydroxyl group is formed. This hydroxyl group is protected with acid and base stable trichloroethoxycarbonyl group (Troc).

A shorter peptide based chain is synthesized by starting with Troc protected 2-(2'-aminoethoxy)ethanol and coupling this with one lysine and two glycines using standard peptide chemistry.

Two peptide chains which are prepared as described above are coupled together by forming an amide bond between the free carboxylic group of glutaminic acid and the end amino group of the glycine in the shorter peptide. The product which has three branches each having one protected hydroxyl group needs manipulation of the protecting groups before it is compatible with oligonucleotide synthesis.

Once the properly protected spacer is prepared, the first prepared oligonucleotide is coupled with phototriester synthesis with the free primary hydroxyl group (Figure 14). The shortest oligonucleotide is coupled in this stage, whereas the longest 5 oligonucleotide is prepared with automatic synthesizer. The product is not deprotected or detached from the solid support. The synthesis is continued by adding the "trivalent" spacer, which is already coupled with one oligonucleotide. The free secondary hydroxyl group becomes coupled with the 10 oligonucleotide which is still bound with the solid support. Thus the peptide spacer is coupled with two oligonucleotide chains. Dimethoxytrityl protecting group of the third hydroxyl group is removed by acid. The automated oligonucleotide synthesis is continued and the third oligonucleotide chain is 15 constructed. The protecting groups are then removed and the molecule is detached from the solid support.

D. Assembly of Supramolecule from Components

The hybridization is performed preferably in a aqueous 20 medium containing various additives. Additives include, but are not limited to buffer, detergent (0.1 % to 1 %) , salts (e.g., sodium chloride, sodium citrate from 0.01 to 0.2 M), polyvinylpyrrolidine, carrier nucleic acid, carrier proteins, etc. Organic solvents may be used in conjunction to water, such 25 as alcohols, dimethyl sulfoxide, dimethyl formamide, formamide, acetonitrile, etc. In addition to concentration of the derivatized oligonucleotides, the temperature can be used to control the hybridization. The optimum temperature for hybridization is 20 °C below the melting point of the 30 oligonucleotide. This means that the preferred temperature for hybridizing 30-mers is typically 40 - 60 °C. For shorter oligonucleotides the temperature is lower and for longer oligonucleotides it is higher. Oligonucleotides containing large portion of cytidine and guanine have higher melting point 35 than the oligonucleotides containing a lot of adenine and thymidine. Detailed formulae for calculating the melting temperature of double stranded nucleic acids are well known to

the person of ordinary skill in the art. Additionally, melting temperature may readily be calculated using empirical methods.

5. Example of Antibody-multienzyme Supramolecule

Two current main strategies for drug development for HIV are finding of reverse transcriptase and HIV protease inhibitors. All four approved AIDS drugs are reverse transcriptase inhibitors. HIV protease inhibitors are also promising as drugs, but the rapid mutation of the viral protease has so far been overwhelming obstacle for the development of a commercial drug.

Embodiments of the supramolecules of the invention that comprise an HIV-antibody and several digestive enzymes can destroy the virus particle itself. Antibodies have earlier been conjugated with enzymes for drug use (C. Bode, M. S. Runge and E. Haber in "The Year in Immunology 1989-1990". Molecules and Cells of Immunity (J. M. Cruse and R. E. Lewis, Eds.) Vol. 6, Karger Publishing, Basel, 1990). Typically these antibody-enzyme complexes are used to produce active drugs from prodrugs. This embodiment of the invention is particularly advantageous if the drug of interest is highly toxic at therapeutic levels. For example, the drug against cancer can be produced on the surface of the cancer cell and cancer cells are subjected to higher concentration of this drug than other cells. Several antibodies specific for cancer cells are known. Enzymes are targeted to degrade the plasma membrane of the cancer cell and include lipases, proteases and glycosidases.

One strategy is to couple lipid and RNA degrading enzymes to an HIV specific antibody. Although a virus does not have its own metabolism to serve as a drug target, a virus is unable to heal itself, if part of the virus is destroyed by externally added catabolic enzymes.

In order for these enzymes to have operational freedom, the spacer between the antibody and the enzyme must be of sufficient, e.g., e.g., on the order of 10 nm. In this case virtually the whole surface of the virus is covered. In order to avoid allergic reactions this spacer must be fully

biocompatible, preferably a normal biological component. In addition it should have some rigidity to allow structures in which enzymes and antibodies do not interfere with each other. Because these antibody-enzyme complexes can be complicated
5 structures, a self-assembly would be ideal. Oligonucleotides fulfill all these requirements. Further requirement is that joints connecting enzymes and antibody with oligonucleotides are as small as possible to suppress immunoreaction. These drugs are supramolecular drugs, i.e., noncovalent interactions are
10 important structural factors. Especially complementary hydrogen bonding of oligonucleotides is essential for the assembly and structural integrity.

In Figure 3A is a schematic representation of one possible supramolecule demonstrating this principle. Antibody is in
15 central position and four different enzymes: phospholipase A₂, lipase, cholesterol esterase and ribonuclease A. Phospholipase A₂ can be supplemented or completely replaced by another phospholipase such as phospholipase C. One extra single stranded oligonucleotide is attached with the antibody. This
20 oligonucleotide is complementary with viral RNA and binds viral RNA when virus is disintegrated.

Many viruses, including HIV-1, are covered by a lipid bilayer which it takes from the host cell when it is formed. The bilayer contains phospholipids, triglycerides and
25 cholesterol esters. Accordingly three enzymes specific for these classes of compounds are used to digest the viral lipid bilayer. When the bilayer is hydrolyzed, fatty acids and lysolipids are formed. These digestion products are soluble in blood plasma and may be bound by albumin, which is a scavenger
30 protein to remove free fatty acids and lysolipids. When the protein core of the virus is exposed to plasma it is to be expected that the protein dissolves spontaneously and RNA is released. This process happens when the virus is internalized into a cell. The lipid bilayer fuses with the plasma membrane
35 of the cell and virus becomes unstable and dissolves into the cytoplasm of the cell. No specific endocytosis mechanism has been observed for HIV. In essence our idea is to induce the

dissolution of the virus outside the cell and destroy viral RNA when it is released. In order to promote the breakdown of RNA a short complementary oligonucleotide is attached with the antibody and also ribonuclease A is part of the enzyme palette.

5 Proteinases are not included among the enzymes in our first design, because it is feasible to suppose that the protein effector of the HIV is unstable when exposed. If opposite turns out to be true, it is possible to include some proteinases. However, blood contains inhibitors against many proteinases,

10 especially if proteinases are nonspecific. Some specific endopeptidases as well as carboxypeptidases and aminopeptidases can be used, because they are not inhibited.

A similar strategy can be used for cancer therapy and to remove 'plaque' from blood vessels, e.g., to treat

15 atherosclerosis. In each case antibody must be replaced with another antibody or other recognition molecule, which is specific for the target. Also enzymatic composition must be adjusted for each application.

The antibody-multienzyme supramolecule is assembled from

20 oligonucleotide-enzyme conjugates and branched oligonucleotides according to Figure 3(B). Figure 3(C) depicts two simplified supramolecules, which together can carry the same enzymes as the supramolecule in Figure 3(A).

An important consideration in the synthesis is the

25 incorporation of amino or thiol functionalities into a desired point of the oligonucleotide during automated synthesis. Phosphoramidite synthesis is described in 5.1-5.7. Their use in oligonucleotide synthesis is straightforward. By using amino and thiol specific cross-linking agents, the synthesis of

30 branched oligonucleotides is also easily accomplished. The oligonucleotide strands are by A and B and their complementary oligonucleotides are denoted by corresponding underlined letters. Enzymes are attached into either 3' or 5'-terminus of the oligonucleotide, which contains an amino group. This kind

35 of coupling of oligonucleotides and proteins is a standard practice in biochemical conjugation. Antibody is attached into

the center of a oligonucleotide chain containing an aliphatic amino group in that position.

After molecular building blocks are synthesized, the final step is a self-assembly of a supramolecule. This relies on the pairwise complementarity of the oligonucleotide strands in the components, which are designed to bind together. In principle, the components contain the complete information of the structure of the final supramolecule and a simple mixing of the component molecules will produce the wanted product. However, in order to make certain that the assembly proceeds as designed, the stepwise process is to be preferred.

Preparation of the antibody-multienzyme supramolecule

Abbreviations: Aminopropanol, AP; 2-Cyanoethyl N,N-diisopropylchloro phosphoramidite, CEDIPCPA; Dichloromethane, DCM; Di-isopropyl ethyl amine, DIPEA; Fluorenylmethoxycarbonyl, Fmoc; Fluorenylmethoxycarbonylchloride, FMOCCL; Methanol, MeOH; Monomethoxytrityl, MMT; Monomethoxytritylchloride, MMTCL; Serinol, SER; Tetrahydrofuran, THF; Triethylamine, TEA.

20

5.1. N-Monomethoxytrityl Aminopropanol (MMT-AP)

MMTCL (1.54 g) in 10 ml of DCM was added to a solution of AP (1.5 ml) in 5 ml of DCM. The reaction mixture was 24 h at + 4 °C. 10 ml of DCM was added and the mixture was washed twice with 10 ml of 5 % NaHCO₃ and with 5 ml of water. The DCM phase was dried with solid NaHCO₃. The solution was concentrated into 5 ml in vacuo and applied to 40 g silica column, which was eluted with 300 ml of DCM/TEA 200:1, 300 ml of DCM/EtOAc/TEA 200:2:1 and 200 ml of DCM/EtOAc/TEA 100:2:1. 1.32 g of pure MMT-AP was obtained.

35

5.2. N-Monomethoxytrityl aminopropyl cyanoethyl N,N-diisopropylphosphoramidite (MMT-AP-CEDIPPA)

To a solution of MMT-AP (0.42 g) in 8 ml of DCM was added 475 μ l of EDIPA and 0.30 g of CEDIPCPA in 2.5 ml of DCM. After 10 min the reaction mixture was applied directly to 10 g silica column. The column was eluted with DCM/EtOAc/EDIPA 98:1:1. Fractions of 6 ml were collected. The product was in fractions 3 and 4. The yield was 0.44 g.

This product was used in oligonucleotide synthesis.

10

5.3. N-Fluorenylmethoxycarbonyl aminopropanol (Fmoc-AP)

FMOCCl (1.55 g) in 10 ml of THF was added into a solution of 0.90 g of AP in 40 ml of water. After 30 min stirring the reaction mixture was extracted with 20 ml of DCM. The DCM solution was washed twice with 10 ml of water and dried with MgSO_4 . The solvent was removed with a rotary evaporator and the residue was dissolved into 14 ml of EtOH and 14 ml of water was added. The small precipitate was filtered off and the solution was put into a refrigerator. After 20 h the precipitate was separated by filtration. The yield was 1.22 g.

20

5.4. N-Fluorenylmethoxycarbonyl aminopropyl cyanoethyl N,N-diisopropylphosphor-amidite (Fmoc-AP-CEDIPPA)

Fmoc derivative was done exactly as MMT analog in Example 2 using 0.30 g Fmoc-AP. Also purification was done similarly. The product was in fractions 3-8. Fractions 3-7 contained 0.41 g product.

25

This product was used in oligonucleotide synthesis.

5.5. N-Fluorenylmethoxycarbonyl serinol (Fmoc-SER)

30

FMOCCl (1.55 g) in 10 ml of THF was added into a solution of 0.54 g of SER in 30 ml of water and 8 ml of 1.5-M Na_2CO_3 . After 30 min stirring the reaction mixture was extracted with 20 ml of EtOAc. The EtOAc solution was washed twice with 10 ml of water and dried with MgSO_4 . The solvent was removed with a rotary evaporator and the residue was dissolved into a mixture

35

of 5 ml of EtOH and 30 ml of DCM. The product crystallized in +4 °C. Yield was 1.12 g.

5.6. N-Fluorenylmethoxycarbonyl O-dimethoxytriphenyl serinol (FMOC-DMT-SER)

5 FMOC-SER (1.12 g) was dissolved into 6 ml of pyridine and 0.68 g of solid DMTrCl was added. The reaction mixture was put into +4 °C. After 20 h 20 ml of water was added and the oily layer was washed with 5 ml of water and dissolved into 10 ml of
10 EtOAc and the solvent was removed in vacuo. The residue (1.76 g) was fractionated in 28 g silica column, which was eluted with DCM/EtOAc/MeOH/TIPEA 98:1;0.2:0.5 and 96:4:1:0.5. Yield of pure product was 0.72 g.

15 5.7. N-Fluorenylmethoxycarbonyl O-dimethoxytriphenyl serinyl cyanoethyl N,N-diisopropylphosphoramidite (FMOC-DMT-SER-CEDIPPA)

FMOC-DMT-SER derivative was produced essentially as described for the phosphoramidite in Example 5.2 using 0.65 g FMOC-AP. The product was purified similarly. The desired
20 reaction product was found in fractions 4-9. Fractions 5-8 contained 0.82 g product. TFMOC-DMT-SER may also be synthesized by first protecting serinol with DMT and then with FMOC. This variation allows also acylation of the amino group of serinol with carboxylic acid carrying various other functionalities,
25 such as protected amino or thiol groups and biotin.

The desired product was used in automated synthesis to introduce aliphatic amino group in the position of 20 in a 51-mer.

30 5.8. Automated Synthesis of Oligonucleotides

The following oligonucleotides were synthesized by automated synthesis:

35 A 3'TGGAGATGGGGCACCATGCTX5'
(SEQ ID NO:1)

B 3'AGCATGGTGCCCCATCTCCAYAGTCACAGCACAGCACTAATAACAAGAA5'
(SEQ ID NO:2)

- C 3' TYTTTCTTGTTATTAGTGCTGTGCTGTGACT5'
 (SEQ ID NO:3)
- D 3' GTGATAGGAGTTGATTACAGTCCTX5'
 (SEQ ID NO:4)
- 5 E 3' AGGACTGTAATCAACTCCTATCACYATCAGAAGAGTGAGACGGTGGGAT5'
 (SEQ ID NO:5)
- F 3' TYATCCCACCGTCTCACTCTTCTGAT5'
 (SEQ ID NO:6)
- G 3' TAGACTCAGCGCAATCGTGAAGCTX5'
 (SEQ ID NO:7)
- 10 H 3' AGCTTCACGATTGCGCTGAGTCTAYGATTCTCGGCTCGTTCGAAGTGTC5'
 (SEQ ID NO:8)
- I 3' TYGACACTTCGAACGAGCCGAGAATC5'
 (SEQ ID NO:9)
- 15 X represents MMT-AP-CEDIPPA (5.2) and Y represents FMOC-
DMT-SER-CEDIPPA (5.7). Analogous amidites may also be to
introduce aliphatic amino groups.

5.9. Purification of Monoclonal Antibody

- 20 Anti gp41/160 (antibody IAM3D6) supernatant had a
concentration of 315 mg/l. It was purified in 160 ml portions
in Protein A Sepharose Fast Flow 5 ml column. The supernatant
was buffered with 40 ml of 0.2-M Na₂HPO₄. After feeding the
supernatant into the column, the column was washed with 120 ml
25 of 0.1-M Na₂HPO₄. The antibody was eluted off the column with
0.1-M citric acid and neutralized immediately with 3-M KOH. The
antibody solutions were stored at -18 °C.

5.10. Acetylated Protein A Sepharose Gel

- 30 Protein A Sepharose was packed into 1.5 ml column. It was
saturated by eluting with a solution containing 50 mg of
monoclonal antibody (Anti gp41/160 IAM3D6). The column was
washed with 0.1-M Na₂HPO₄ buffer (15 ml) and eluted 10 ml 1 mM
acetyl N-hydroxy succinimide solution in DMF/water 1:9. The
35 antibody was removed by 0.1-M citric acid. The acetylated
Protein A Sepharose was used to couple antibody with nucleotides
and in the final assembly of the supramolecule.

5.11. Coupling of Oligonucleotide with Antibody

A solution of antibody (40 mg/25 ml water) was eluted through the column containing 1.5 ml acetylated Protein A Sepharose. The Sepharose was washed with 3 ml of 0.1-M Na_2HPO_4 and taken out of the column to perform a bath reaction with derivatized nucleotide.

Oligonucleotide 2 (10 mg, 0.5 μmole), comprising two equal 30-mers bound together by an amino group containing joint, was dissolved into 1 ml of 0.1-M NaHCO_3 and 50 μl of 1-M solution of bis(hydroxysuccinimidyl) glutarate in acetonitrile was added. After one hour the water solution was extracted twice with 1 ml of EtOAc and the solution was dialyzed 2 h against 0.1-M NaHCO_3 . The activated nucleotide was added into a slurry of Sepharose. The mixture was stirred six hours and packed into a column. The antibody coupled to the nucleotide was eluted off the column with 0.1-M citric acid. Antibody-oligonucleotide conjugate was fractionated in a Sephadex G-25 column and antibody connected with oligonucleotide was collected.

5.12. Coupling of Oligonucleotide with Enzymes

Oligonucleotide 1 (20 mg, 2 μmole), which was contained aliphatic amino group at 5'-position was dissolved into 2ml of 0.1-M NaHCO_3 and 400 μl of 1-M solution of bis(hydroxysuccinimidyl) glutarate in acetonitrile was added. After one hour the water solution was extracted twice with 1 ml of EtOAc. The solution was dialyzed 2 h against 0.1-M NaHCO_3 and 0.5 ml aliquots of this solution were added into the following enzyme solutions:

- a. 10 mg phospholipase A_2 in 1 ml of water.
- b. 40 mg lipase in 4 ml of water.
- c. 10 mg ribonuclease in 1 ml of water
- d. 30 mg carboxypeptidase in 3 ml of water

5.13. Assembly of the supramolecule

Antibody connected with oligonucleotide was eluted through a acetylated Protein A Sepharose column (1.5 ml) so that the column was saturated with antibody. The column was thermostable at + 40 °C and phospholipase-oligonucleotide conjugate solution (twice the equivalent amount) was circulated through the column and UV-flow cuvette. When UV-absorption at 280 nm was decreased into half the ribonuclease A-oligonucleotide conjugate was circulated similarly through the column. Generally about two hours was needed for a complete reaction. The supramolecule was eluted off the column by 0.1-M citric acid and neutralized immediately with 1-M KOH. The other supramolecule depicted in Figure 3(C) was prepared similarly.

6. Design Of Supramolecule For Capturing Virus Particles

This example describes the design of a supramolecular assembly that is capable of surrounding a comparatively large particle, e.g, a virus. First, a structure, which is capable of performing the desired function, is designed and the geometrical features are fixed. Then chemical and physical features are chosen based on the application. Hydrophilicity, hydrophobicity, acidity, alkalinity, charge transfer, etc., is mapped onto the structure. This designed structure may be visualized as a single molecule, although in many instances the synthesis of this molecule would be difficult to achieve at a reasonable yield. In such embodiments, supramolecular retrosynthesis is performed, i.e., the structure is broken down into small molecules, which are capable via self-assembly of forming the original structure. The supramolecular assembly produced in this manner is not identical with the molecule represented as a schematic in the figures; however, the important characteristics, i.e., geometry and chemical and physical properties listed above, remain the same. Supramolecular retrosynthesis does not try to retain the original molecular structure intact, but tries to retain all the important chemical and physical properties of the desired structure.

Another retrosynthetic cycle can be performed for the molecules obtained in the previous retrosynthesis to obtain smaller molecular building blocks. Finally, molecules are obtained that can be designed and prepared easily. In the 5 design example given below, there are two retrosynthetic cycles.

Many viruses have an icosahedral shape. Such a virus can be covered by an icosahedral and assembly designed according to this invention. This process is demonstrated stepwise in Figures 5 A-E. Dimensions referenced are taken from HIV (human 10 immunodeficiency virus), but the same principles apply to any virus. In this example, polypeptides and oligonucleotides are used, because synthetic methods are available for their high yield synthesis. As synthetic methods further develop, analogues or completely artificial supermolecular systems can be 15 made using the same design and construction principles offered by the invention.

Each edge of HIV is about 80 nm. In preliminary design we suppose that three amino acid or nucleic acid residues are needed per nanometer. The circles in Figure 4 represent cyclic 20 polypeptides containing enough lysine so that five polylysine chains can be attached. These polylysines are denoted by zig-zag lines in Figure 4. Polylysine should contain about 200 residues in order to cover whole edge. Onto the other end of each polylysine chain is coupled another cyclic peptide that has 25 four nucleotides attached. These nucleotide strands are denoted by a wavy line in Figure 4. Two of these oligonucleotides are the same, for example, oligonucleotide 1 ($n=p=1$, $i=100$). Two others are mutually complementary, but they are bound to the cyclic peptide so that coupling occurs easily between neighbors, 30 but not intramolecularly. Thus, they form a pentagon shaped double helix. Figure 4 shows single stranded oligonucleotide is bound by polylysine. Another molecule is designed using the same principles, but instead of oligonucleotide 1, the single stranded oligonucleotide is now 1. When either of these 35 molecules encounters a virus, which has a negatively charged surface, polylysine is Coulombically associated with the virus. Simultaneously, a negatively charged oligonucleotide (e.g., 1 or

1) is released from the polylysine. When a complementary capping molecule is associated with the virus, the complementary oligonucleotides (1 and 1) combine to close the cage from which the virus can not escape.

5 The molecules shown in Figures 4-5 and described above would be incredibly difficult to synthesize. However, by designing a supramolecular constructed from smaller components, synthesis of a virus capturing molecule is made possible.

Figure 6 demonstrates how the analogous structure for the
10 large molecule in Figure 4 can be prepared using smaller molecules. These smaller molecules are shown separately in Figure 6. Thus, in this approach six different compounds are needed to get the overall structure, which is same as that of the molecule in Figure 4. Four of these are relatively simple,
15 because in each two oligonucleotides are connected to a spacer, which can be polypeptide. Two molecules in the upper part of Figure 4 are still relatively complicated because in both cases five nucleotides are connected to a cyclic spacer, which can be a cyclic peptide. These oligonucleotides are denoted by
20 (3,3,3,3,3) and (1,1,8,7,8). In these notations only free single stranded oligonucleotides are listed. These structures can be synthesized by attaching each type of oligonucleotide needed to a short peptide, for example, pentapeptide Gly-Ala-Ser-Ala-Gly which is otherwise protected but the hydroxyl group of serine is
25 free. Nucleotide is connected with this hydroxyl group using normal phosphate coupling. Then, using peptide synthesis methods, these pentapeptides connected to a specific oligonucleotides are coupled in a desired order. Closing the cycle makes the molecule more symmetric, but is not essential
30 for the supramolecular assembly or the function of this assembly in most cases.

There is a further possibility of assembly of cyclic structures containing five oligonucleotide chains by using the general principles of this application. This second step of
35 supramolecular retrosynthesis is demonstrated in Figure 8. Both of these cyclic structures can be assembled from five smaller molecules. For (3,3,3,3,3) these molecules are twice (3,1,2) and

once (3,2,4), (3,2,4), (3,1,2) and for (1,1,8,7,8) these are (2,1,6), (6,1,5), (5,8,4), (4,7,3), (3,8,2). In these notations it is immediately clear that molecule (2,1,6) combines with the molecule (6,1,5), because complementary nucleotides are written last and first, respectively. Looking at the whole sequence of five molecules indicates that the notation starts with nucleotide 2 and ends with 2. This means that these ends will bind together and form a pentagon. After assembly, a supramolecule is obtained which has the same overall shape as two molecules in upper part of Figure 7. These supramolecules are still denoted by listing only their single stranded oligonucleotides, because this is important for further assembly and is sufficient for purposes of this application. The symbols are (3,3,3,3,3) and (1,1,8,7,8). These supramolecules also function similarly in further assembly of the structure, which has the same shape as the molecule in Figure 4. This demonstrates that almost any structure can ultimately be created from molecules which has a spacer or a molecular moiety having an active role in the final assembly connected to two or three oligonucleotides. The spacer can be a very small molecule or it can be a large molecule. The spacer can actually be a DNA strand.

Supramolecular assemblies are preferably prepared in an aqueous environment, although some embodiments may be assembled in organic solvents. When effector molecules are lipophilic, the Langmuir-Blodgett technique may be utilized. Stepwise assembly is often advantageous. For example, the cyclic structures (3,3,3,3,3) and (1,1,8,7,8) in Figure 8 are assembled separately. These two structures can be stabilized internally by cross-linking their double helices. This cross-linking can be performed in a highly selective manner. By cross-linking, both of these supramolecular assemblies become covalent molecules. Cross-linking is not essential, but can be advantageous, because it increases thermal stability. After first assembly and possible cross-linking, the product can be purified. Purification as well as cross-linking is to be

recommended, if the same oligonucleotide is used in several different places.

During the second assembly step (3,4) and (4,7) (see Figure 6) are added to (3,3,3,3,3) to give (7,7,7,7,7). In the third assembly step the product (7,7,7,7,7) and (1,1,8,2,8) are combined to form $10^*(1,8)$. The fourth assembly step is the formation of a pentagon by adding (8,9) and (8,9) to give $10^*(1)$. The fifth and final assembly step is adding single stranded oligonucleotide 1, and the end product is $10^*(1)$.
10 After each step, cross-linking or purification or both can be performed depending on the final requirements regarding quality of the product. The complementary supramolecule $10^*(1)$ is prepared similarly.

If necessary, the cage surrounding the virus can be made
15 more dense using the principles of this application. The number of molecules needed is then correspondingly larger.

DNA double helix is thermally unstable and cross-linking may be required for stability. One possible approach is shown in Figure 12. The last amino acid residue in the spacer is
20 lysine and a complementary DNA strand contains an alkylating group, which binds preferentially with the amino group of lysine, because it is the most nucleophilic of the functional groups in this assembly. Thus, perfect chemical control can be maintained also in the cross-linking process, although this is
25 not always necessary and more random cross-linking methods can be used. Incorporating photoactivatable groups, like azido adenosine or bromo- or iodo uridine, into oligonucleotide chains allows photochemical cross-linking, which is site specific. Also use the 3-thioribose in oligonucleotide and cysteine in the
30 peptide spacer allows formation of disulphide bridges.

7. Preparation of Components for Liposomes

Many naturally occurring proteins may be used as such without any modification. Because in some cases these liposomes
35 are used *in vivo* in humans, the human proteins should be used to minimize unwanted host immune responses. Human proteins can be produced in transgenic plants or animals. The amino acid

sequence of proteins may also be altered through well-known genetic engineering techniques to produce mutated proteins having the desired biological functions of corresponding naturally occurring protein, but adapted to coupling to lipid
5 molecules. For example, addition of a cysteine residue, either through substitution or inserting, will add a free thiol group for coupling to a lipid molecule. Moreover the location of this cysteine can be deliberately chosen. It should be located so that conjugation does not disturb the activity of the protein.
10 Normally this means that the cysteine must be as far as possible from the active site.

In order to covalently couple a lipid with an effector molecule, the lipid must contain a functional group which has
15 high enough reactivity with heterobifunctional cross-linker. Phosphatidylethanol-amines (PE) have free aliphatic amino group and can be used directly for conjugation. The conjugation of phosphatidylethanolamines (PE) and serines (PS) with proteins is well known in the literature (Egger et. al. Biochem. Biophys.
20 Acta 1104 (1992) 45-54). Especially, if the protein contains a thiol group, a chemically well defined conjugate may be easily prepared. Several spacers are commercially available to couple two molecules, which contain an amino and a thiol group. If a protein does not contain an a thiol group, an aliphatic amino
25 group, may be used for conjugation. Thiol groups are preferred for conjugation sites in proteins as compared with amino groups because of the lower abundance of thiol groups in most proteins.

Proteins can also be conjugated with lipids after the
30 preparation of the liposomes so as to position the proteins on the outer surface of the liposome.

Phospholipids having unnatural stereochemistry may be prepared as described in the literature, for example see,
35 (J.A. Virtanen et. al., Chem. Phys. Lipids 27 (1980) 185). D-Mannitol is tritylated to yield 1,6-ditrityl-D-mannitol, which is oxidized with lead tetra-acetate and the product reduced

with sodium borohydride to yield 1-trityl-sn-glycerol (1-TrG). Diacylglycerols and phospholipids can be prepared starting from 1-TrG by standard methods. For the liposomes described in this application ether bond is preferred at least in the
5 primary hydroxyl group, because it is resistant against lipase. in sn-hydroxyl ester bond can be used, because unnatural stereochemistry will make this bond stable against phospholipase A2. Instead of glycerophospholipids sphingolipids or completely artificial lipids, which are
10 resistant against these enzymes, can be used.

8. Examples of effector molecule combinations for specific diseases

Binding molecule-multienzyme complexes may be designed so
15 as to treat diseases caused by specific pathogenic agents. Examples of effector molecule combinations are provided below in table 3 so that from each of the three columns (titled:antibodies, enzymes and peptides) suitable effector molecules are chosen. Binding molecule-multienzyme complexes,
20 as described in this example, may contain one or several antibodies, which can be different or the same as one another. The binding molecule-multienzyme complexes may comprise 3-8 different enzymes as therapeutic effector molecules, which are typically selected so that 1-3 of the enzymes degrade lipids,
25 0-3 of the enzymes degrade proteins, 1-2 of the enzymes degrade nucleic acids and 0-3 of the enzymes degrade carbohydrates. Chemotactic peptides may be used as effector molecules enhance the natural defence mechanisms of the organism for treatment. Platelet aggregation inhibitors may
30 also be used as effector molecules to prevent bacterial adhesion to human cells. Additionally, components such as coenzymes, cholic acids, polyamines and metal chelates can be included. These additional compounds may be used to activate enzymes or facilitate accessibility to the binding target so
35 that enzymatic therapeutic effector molecules may more readily degrade the target. The binding molecule-multienzyme complexes may be also loaded into another assembly so that

enzymes are activated at the target site. For example, instead of active enzymes, proenzymes may be used. When proenzymes are employed as therapeutic effector molecules. In this case a small amount of activating enzyme is brought into the target.

Table 3

VIRUSES

10	Antibodies:	Enzymes:	Peptides:
	Anti gp41 (HIV)	Phospholipase A ₂	Chemotactic
	Anti gp120 (HIV)	Phospholipase C	peptides, e.g.,
		Lipase	N-Formyl-Met
	Anti hepatitis B	Cholesterol esterase	Leu-Phe
15		Cholesterol oxidase	
		Aminopeptidase	
		Endoproteinase Arg-C	
		Endoproteinase Asp-N	
20		Endoproteinase Lys-C	
		Carboxypeptidase A	
		Carboxypeptidase B	
		Chymotrypsin	
25		Ribonuclease A	
		Ribonuclease B	
		Ribonuclease C	

BACTERIA, PROTOZOA AND FUNGI

30	Antibodies:	Enzymes:	Peptides:
	Anti TB	α -Amylase	Chemotactic
	(Mycobacterium tuberculosis)	β -Amylase	peptides, e.g.,
		Galactosidase	N-Formyl-Met-
		Galactose oxidase	Leu-Phe
35	Anti Syphilis	α -Mannosidase	

	(Treponema pallidum)	β -Mannosidase	Platelet aggregation inhibitors,
	Anti Cholera	Phospholipase A ₂	e.g., Arg-Gly-Asp-
5	(Vibrio Cholerae)	Phospholipase C	Ser
		Lipase	
		Cholesterol esterase	
		Lysozyme	
		Lactoferrin	
10	Anti plasmodium	Aminopeptidase	
		Endoproteinase Arg-C	
		Endoproteinase Asp-N	
		Endoproteinase Lys-C	
15		Carboxypeptidase A	
		Carboxypeptidase B	
		Chymotrypsin	
		<u>CANCER</u>	
20	Antibodies:	Enzymes:	Peptides:
	HUMAbSK1	α -Amylase	Chemotactic
		β -Amylase	peptides, e.g.,
	Humanized	Galactosidase	N-Formyl-Met-
	antimucin	Galactose oxidase	Leu-Phe
25		α -Mannosidase	
		β -Mannosidase	Platelet aggregation inhibitors,
		Phospholipase A ₂	e.g., Arg-Gly-Asp-
		Phospholipase C	Ser
30		Lipase	
		Cholesterol esterase	
		Aminopeptidase	
		Endoproteinase Arg-C	
35		Endoproteinase Asp-N	
		Endoproteinase Lys-C	
		Carboxypeptidase A	

Carboxypeptidase B
Chymotrypsin

Alkaline phosphatase

5 Polyphenol oxidase

ATHEROSCLEROSIS

Antibodies:	Enzymes:	Peptides:
Murine/Human	Phospholipase A ₂	Chemotactic
10 IgG1Z2D3	Phospholipase C	peptides, e.g.,
	Lipase	N-Formyl-Met-
	Cholesterol esterase	Leu-Phe
	Cholesterol oxidase	
15	Aminopeptidase	Platelet aggre-
	Endoproteinase Arg-C	gation inhibitors,
	Endoproteinase Asp-N	e.g., Arg-Gly-Asp-
	Endoproteinase Lys-C	Ser
	Carboxypeptidase A	
20	Carboxypeptidase B	
	Chymotrypsin	
	Collagenase	

9. Examples of liposome formulation adapted for the
25 treatment of HIV infections

ABBREVIATIONS

Buffers:

30 BA: 10mM NaH₂PO₄ + 0.1 M NaCl

BB: 10mM Na₂HPO₄ + 0.1 M NaCl

35 BD: 0.25 M dithiotreitol in BA

MOPS: 3-(N-morpholino) propanesulfonic acid

PBS: Standard phosphate saline buffer

5 Spacers and conjugation agents:

MIPEG: Maleimide polyethyleneglycol

BisMIPEG: Bismaleimide polyethyleneglycol

10

SPDP: 3-(2-Pyridyldithio) propionic acid N-hydroxysuccinimide ester

15 Enzymes:

PLase: Phospholipase A₂

RNAse: Ribonuclease A

20

Phospholipids:

25 OSPC: 1-Octadecyl-2-stearoyl phosphatidylcholine

OSPE: 1-Octadecyl-2-stearoyl phosphatidylethanolamine

OSPE-PDP: N-[3-(2-Pyridyldithio) propionyl]-OSPE

30

35

ANTIBODY REDUCTION AND MIPEG-Ab

35 mg (0.25 μ moles) Ab in 5 ml of PBS is dialyzed 3 x 2 h
5 against 19:1 BA/BB at RT.

Into this solution is added 0.5 ml of 0.1 M thioethanolamine
in 9:1 BA/BB.

10 Kept 1 h at 37°C.

Dialyzed under nitrogen 3 x 2 h against 9:1 BA/BB, which is
purged with nitrogen.

15 A 50 μ l sample diluted with 2 ml of BA and UV-spectrum is
measured between 240-400 nm. The absorbance at 280 nm should
be about 0.21. 50 μ l of SPDP is added. Absorbance is
measured at 343 nm immediately after the addition and after 10
min. The increase in the A_{343} should be about 0.024.

20

160 μ l of 50 mM Bis MIPEG in EtOH (8 μ moles) is added and let
stand 2 h under nitrogen at RT.

Dialyzed 3 x 2 h against 9:1 BA/BB.

25

The A_{280} should be about 0.20 and A_{305} should be about 0.025.

The MIPEG-PLase solution is transferred into a bottle.

30

THIOL-CD4 AND MIPEG CD4

35

54 mg (1 μ mol) of sCD4 in 6 ml is dialyzed 3 x 2 h against BB in dialysis tube (MWCO 15,000).

2.5 μ moles of SPDP in 250 μ l of EtOH added and allowed to stand 3 h at RT.

Dialyzed 2 x 2 h against BA.

A 200 μ l sample diluted with 1.8 ml of BA and UV-spectrum is measured between 240-400 nm. The absorbance at 280 nm should be about 0.64. 50 μ l of BD is added. Absorbance is measured at 343 nm immediately after the addition and after 10 min. The increase in the A_{343} should be about 0.10.

15 1 ml of BD is added into the main mixture and let stand 30 min at RT.

Dialyzed 3 x 2 h against BA. Each dialysis is performed against nitrogen purged BA and under nitrogen.

20

A 200 μ l sample diluted with 1.8 ml of BA and UV-spectrum is measured between 240-400 nm. The absorbance at 280 nm should be about 0.48. 50 μ l of 10 mM SPDP in EtOH is added. Absorbance is measured at 343 nm immediately after the addition and after 10 min. The increase in the A_{343} should be about 0.06.

Meanwhile, 6 μ moles of BisMIPEG is dried under nitrogen blow.

30 CD4 solution is added onto BisMIPEG. The solution is purged gently with nitrogen and let stand under vacuum 2 h.

The reaction mixture is transferred into the dialysis tube (MWCO 15,000) and dialyzed against nitrogen purged BA and under nitrogen overnight at +4°C.

35

25 μ l 20 mM MIPA solution is added and dialysis is continued 2
x 2 h against BA under ambient.

The MIPEG-CD4 solution is transferred into a bottle.

5

MIPEG-PLase

10

50 mg (4 μ moles) PLase is dissolved into 4 ml of 15 mM MOPS and
dialyzed 2 h against 15 mM MOPS at RT.

Into this solution is added:

15

40 MG (20 μ moles) of MIPEG

56 mg (160 μ moles) of hexadecylphosphorylcholine (HDPC)

and mixed gently until HDPC is completely dissolved (as
20 micelles).

As a solid are added:

7.2 mg (60 μ moles) of N-hydroxysuccinimide (NHS)

25 57 mg (300 μ moles) of 1-Ethyl-3-(3-
dimethylaminopropyl)carbodiimide methiodide (EDC)

and mixed shortly and let stand at RT 6 h or more (up to 16
h).

30

Dialyzed 3 x 2 h against 9:1 BA/BB.

A 100 μ l sample diluted with 1.9 ml of BA and UV-spectrum is
measured between 240-400 nm. The absorbance at 280 nm should
35 be about 0.24.

The MIPEG-PLase solution is transferred into a bottle.

5

MIPEG-Lipase

50 mg (=1 μ mole) lipase is dissolved into 4 ml of BB and dialyzed 2 h against BA at RT.

10 A 200 μ l sample diluted with 1.8 ml of BA and UV-spectrum is measured between 240-400 nm. The absorbance at 280 nm should be about 0.27.

Meanwhile 5 μ moles of BisMIPEG is dried under nitrogen blow.

15

Lipase solution is added onto BisMIPEG. The solution is purged gently with nitrogen and let stand under vacuum 2 h.

The reaction mixture is transferred into the dialysis tube
20 (MWCO 15,000) and dialyzed against nitrogen purged BA and under nitrogen overnight at +4°C.

25 μ l 20 mM MIPA solution is added and dialysis is continued 2 x 2 h against BA under ambient.

25

The MIPEG-Lipase solution is transferred into a bottle.

30

THIOL-RNase AND MIPEG-RNase

35 55 MG (4 μ moles) RNase is dissolved into 4 ml of BB and dialyzed 2 h against BB at RT.

8 μ moles of SPDP in 0.8 ml of EtOH added and allowed to react 2 h at RT.

Dialyzed 2 x 2 h against BA.

5

A 200 μ l sample diluted with 1.8 ml of BA and UV-spectrum is measured between 240-400 nm. The absorbance at 280 nm should be about 0.16. 50 μ l of 10 mM SPDP in EtOH is added.

Absorbance is measured at 343 nm immediately after the addition and after 10 min and 20 min. The increase in the A_{343} should be about 0.09 in 10 min and 0.15 in 20 min.

Meanwhile 18 μ moles of BisMIPEG is dried under nitrogen blow.

15 CD4 solution is added onto BisMIPEG. The solution is purged gently with nitrogen and let stand under vacuum 2 h.

The reaction mixture is transferred into the dialysis tube (MWC0 15,000) and dialyzed against nitrogen purged BA and under nitrogen overnight at +4°C.

20 25 μ l 20 mM MIPA solution is added and dialysis is continued 2 x 2 h against BA under ambient.

25 The MIPEG-RNase solution is transferred into a bottle.

30

PREPARATION OF LIPOSOMES

35 The following components are dissolved into 10 ml of 5:5:0.1 dichloromethane/ethanol/water:

47 mg	(66 μ moles)	OSPC
47 mg	(66 μ moles)	Sphingomyelin
8 mg	(12 μ moles)	OSPE
3 mg	(3 μ moles)	OSPE-PDP
5 58 mg	(150 μ moles)	Cholesterol

The mixture is heated at 50°C in a wide-neck flask under nitrogen blow, while simultaneously adding dropwise dichloromethane to prevent crystallization of any of the
10 components. When almost all solvent has evaporated, the dichloromethane addition is stopped and nitrogen blow is continued 20 min at 50°C. Onto semisolid residue is added 27 ml of warm (50°C) 9:1 BA/BB.

15 The mixture is sonified 2 min with Branson tip sonifier using output of 4.

A 40 μ l sample diluted with 2 ml of BA and UV-spectrum is measured between 240-400nm. The absorbance at 280 nm should
20 be about 0.40.

The Liposome solution is transferred into a bottle.

25

30

35

CONJUGATION OF THE PROTEINS WITH THE LIPOSOME

2 ml of the liposome solution is filtered through 0.2 μ m
5 filter (using high pressure filtration will minimize the loss
of liposomes). Into the filtered liposome solution is added
0.2 ml of BD. Let stand 30 min at RT and dialyzed under
nitrogen 3 x 2 h against 9:1 BA/BB.

10 Into the dialyzed liposome solution is added:

0.35 ml MIPEG-Ab

0.25 ml MIPEG-CD4

0.1 ml MIPEG-PLase

15 0.2 ml MIPEG-Lipase

0.1 ml MIPEG_RNase

Let stand under nitrogen 4 h at RT.

20 Fractionated in Bio-Gel A5m column (1 cm x 30 cm) eluted with
1 mM 2:3 NaH₂PO₄/Na₂HPO₄ + 0.1 M NaCL. First peak is the
liposomal drug eluted in about 8 ml.

The absorbance at 280 nm should be about 0.40.

25

The Liposomal drug solution is transferred into a bottle.

30

DOSAGE

35 About 90% of the mass of the liposomal drug consists of
phospholipids and the rest of the five proteins. In the
following, the total mass is indicated.

The minimum and maximum doses are estimated to be 9 mg and 300 mg, respectively. The most probable dose is about 50 mg. If the drug cures the patient, it is given every day during about two weeks. If it is not cured, it is given every other day or 5 twice a week continuously.

10

15

20

25

30

35

Incorporation by Reference

All patents, patents applications, and publications cited are incorporated herein by reference.

5 Equivalents

The foregoing written specification is considered to be sufficient to enable one skilled in the art to practice the invention. Indeed, various modifications of the above-described makes for carrying out the invention which are
10 obvious to those skilled in the field of molecular biology, organic chemistry, or related fields are intended to be within the scope of the following claims.

15

20

25

30

35

CLAIMS

What is claimed is:

1. A binding molecule-multienzyme complexes comprising:
a binding effector molecule, and
5 a therapeutic effector molecule,
and a joining component, wherein said component is a
liposome and said binding effector molecule and said
therapeutic effector molecule are functionally attached to the
surface of said liposome.
10
2. The binding molecule-multienzyme complexes
according to Claim 1, wherein said binding effector molecule
is a protein.
- 15 3. The binding molecule-multienzyme complexes
according to Claim 2, wherein said binding effector molecule
is selected from the group consisting of antibodies and
receptor ligands.
- 20 4. The binding molecule-multienzyme complexes
according to Claim 3, wherein said binding effector molecule
is CD4.
5. The binding molecule-multienzyme complexes
25 according to Claim 1, where said therapeutic effector molecule
is a protein.
6. The binding molecule-multienzyme complexes
according to Claim 5, wherein said therapeutic effector
30 molecule is an enzyme.
7. The binding molecule-multienzyme complexes according
to Claim 6, wherein said enzyme is selected from the group
consisting of proteinases, lipase, esterases, and toxins.
35
8. A binding molecule-multienzyme complex adapted for
genetic manipulation complex comprising a binding molecule-

multienzyme complex according to claim 1 and a genetic manipulation complex, said genetic manipulation complex comprising a polynucleotide for genetic manipulation comprising a polynucleotide sequence of interest and a DNA binding protein binding site, a DNA binding protein bound to said DNA protein binding site, motor protein, wherein said motor protein is conjugated to said polynucleotide, and wherein said liposome comprise at least one binding effector conjugated to the surface of said liposome.

10

9. A binding molecule-multienzyme complex adapted for genetic manipulation complex according to Claim 8, wherein said liposome further comprises a phospholipase.

15 10. A binding molecule-multienzyme complex adapted for genetic manipulation complex according to Claim 8, wherein said phospholipase is phospholipase A2.

11. A binding molecule-multienzyme complex adapted for genetic manipulation complex according to Claim 9, wherein a nuclear receptor binding protein is conjugated to said polynucleotide.

12. A binding molecule-multienzyme complex adapted for genetic manipulation complex according to Claim 9, wherein said polynucleotide encodes a therapeutic polypeptide.

13. A method of genetically modifying a cell of interest in a mammal, said method comprising the step of administering a binding molecule-multienzyme complex adapted for genetic manipulation complex according to claim 1.

14. A method of genetically modifying a cell according to Claim 13, wherein said liposome further comprises a phospholipase.

15. A method of genetically modifying a cell according to Claim 13, wherein said phospholipase is phospholipase A2.

5 16. A method of genetically modifying a cell according to Claim 14, wherein a nuclear receptor binding protein is conjugated to said polynucleotide.

17. A method of genetically modifying a cell according to
10 Claim 14, wherein said polynucleotide encodes a therapeutic polypeptide.

15

20

25

30

35




Figure 1




Figure 2 A. Formation of square using four molecules connected by complementary oligonucleotide strands. The numbers refer to Table 1. In oligonucleotides between P-Q, $n=1$ and $p=2$ and in all other nucleotides $n=2$ and $p=1$. The number of repeating units i can be chosen freely, but is same for all nucleotides in this case.




Figure 2 B. Square in Fig. 2 A can be stabilized by diagonals, which have length of 1.41 times the sides.




Figure 2 C. Tetrahedron is formed, if M-Q and P-N connections are of same length as the sides of the square in Fig. 2A.




Figure 3 A. Schematic representation of a supramolecule which is capable selectively binding to a HIV and destroying it.




Figure 3 B. Assembly of the antibody-multienzyme complex.




Figure 3 C. Two supramolecules used to destroy a virus.




Figure 4. The molecule which is capable of forming a cage around a virus when it combines with a complementary molecule.




Figure 5 A. Typical icosahedral virus. The edge of HIV is about 80 nm. Perspective is enlarged so that also a part of the backside is drawn by solid lines.




Figure 5B. A drug molecule of Fig. 4 is approaching the virus.




Figure 5C. Another drug molecule is approaching from below.




Figure 5 D. Two drug molecules have attached onto the virus.




Figure 5E. Two drug molecules bind together by complementary DNA strands.





Figure 6. Construction of the structure represented by molecule in Fig. 4 from smaller molecules. The supramolecule formed from these molecules has same overall shape, but its chemical structure is in many places different. Because of fivefold symmetry axis only part of the assembly is shown.




Figure 7. The molecules, which are needed to construct the supramolecular assembly according to Fig. 6. Symbolism used in this application is shown under the molecules. Numbers refer to Table 1.




Figure 8. The supramolecular assemblies, which give analogous structures than the two molecules shown above and in Fig. 7.




Figure 9. Using triple helix binding in supramolecular assembly. Next step in this assembly is to add $5 * (4, \underline{7})$. This is an alternative to the assembly route in Figs. 6-8.




Figure 10. One possible spacer type for connecting three different oligonucleotides. Peptide part can contain amino acids, which have specific functions. TMT = Trimethoxytrityl, MTr = Methoxytrityl.





Figure 11. An alternative peptide spacer type for connecting three different oligonucleotides.



X = Cl, Br, I

Y = N, S




Figure 12. One possible method for site specific cross-linking. Underlined oligonucleotide is complementary to one already connected to peptide and contains covalently bound cross-linking agent.









Figure 13. Coupling of two derivatized peptide chains to get a branched 'trivalent' spacer.
AAE is 2-(2'-Aminoethoxy) ethanol, PD is propanediol. Other symbols have their standard meaning.




Figure 14. Using spacer which contains one oligonucleotide in the automated oligonucleotide synthesis to obtain 'trivalent' building block for supramolecular assembly.




Figure 15A. Schematic representation of supramolecular system, comprising of two supramolecules, which is capable of cell selective transfection. Arrow indicates the restriction site.




Figure 15B. Schematic representation of supramolecular system, comprising of two supramolecules, which is capable of cell selective transfection.




Figure 15C. Formation and internalization of a supramolecule.





Fig. 16. Conjugation of a phospholipid and protein. These conjugates are used to form a liposome.



Y	Antibody
U	Urokinase
L	Lipase
A	Phospholipase A ₂
C	Cholesterol esterase

Fig. 17. Liposome containing antibody and urokinase lipase, phospholipase A₂ and cholesterol esterase which will hydrolyze the atherosclerotic plaque.




Fig. 18. Antibody-CD4-multienzyme liposome and its binding HIV-1. CD4 will bind to viral coat protein gp120 and antibody to gp41. Enzymes have affinity to viral lipid bilayer and will diffuse between the liposome and virus.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGES CUT OFF AT TOP, BOTTOM OR SIDES**
- ☒ **FADED TEXT OR DRAWING**
- ☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE (S) OR EXHIBIT (S) SUBMITTED ARE POOR**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox